

10/580,011

=> file caplus

FILE 'CAPLUS' ENTERED AT 15:26:31 ON 31 JAN 2008

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2008 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 31 Jan 2008 VOL 148 ISS 5


FILE LAST UPDATED: 30 Jan 2008 (20080130/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply. They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> d que

L1 STR



G1 H,Ak

G2 Cb,Ak

Structure attributes must be viewed using STN Express query preparation.

L3 550 SEA FILE=REGISTRY SSS FUL L1

L4 57 SEA FILE=CAPLUS L3

=> d l4 1-57 ibib abs hitstr

L4 ANSWER 1 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:1168693 CAPLUS

DOCUMENT NUMBER: 148:17977

TITLE: pH indicator titration: a novel fast pKa determination method

AUTHOR(S): Kong, Xiaole; Zhou, Tao; Liu, Zudong; Hider, Robert C.

CORPORATE SOURCE: Division of Pharmaceutical Science King's College London, London, SE1 9NH, UK

SOURCE: Journal of Pharmaceutical Sciences (2007), 96(10), 2777-2783

CODEN: JPMSAE; ISSN: 0022-3549

PUBLISHER: Wiley-Liss, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

10/580,011

AB This study describes a fast spectrophotometric titration method for apparent ionization constant (pKa) determination. In this method, a Universal pH indicator is


utilized instead of the conventional pH electrode. An auto-burette is set to add HCl at a constant rate to a vigorously stirred 1 cm UV cuvette which contains sample and indicator solution. A spectrophotometer continuously records the spectra. Acquired spectral data are processed by calculating the pH from the indicator spectra in the visible region and extracting sample spectra from the UV region. Five compds. possessing pKa values in the range 2-10 were investigated. These results differed from measurements by conventional spectrophotometric titration by ± 0.05 to ± 0.10 log unit.

IT 243987-44-2 349141-34-0

RL: ANT (Analyte); PRP (Properties); ANST (Analytical study)
(pH indicator titration for fast pKa determination)


RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



RN 349141-34-0 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,6-dimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 2 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2007:565038 CAPLUS

DOCUMENT NUMBER: 147:9803

TITLE: Preparation of hydroxydihydropyridinones, hydroxydihydropyridinethiones, hydroxypyranones, and hydroxypyranthiones as metalloprotein inhibitors

INVENTOR(S): Puerta, David T.; Cohen, Seth M.; Lewis, Jana A.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 40pp., Cont.-in-part of Appl. No. PCT/US2005/014747.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English





FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2007117848	A1	20070524	US 2006-554475	20061030
WO 2005110399	A2	20051124	WO 2005-US9277	20050321
WO 2005110399	A3	20060615		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2006028523	A2	20060316	WO 2005-US14747	20050428
WO 2006028523	A3	20060601		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
PRIORITY APPLN. INFO.:			US 2004-566882P	P 20040429
			US 2004-576444P	P 20040603
			WO 2005-US9277	A 20050321
			WO 2005-US14747	A2 20050428
			US 2006-826488P	P 20060921

OTHER SOURCE(S):
GI

MARPAT 147:9803



AB The compds. (I), (II), (III), and (IV) [wherein X = O, S; one or two of R1, R2, R4, and R4 is individually a substituent of formula [C6-10 aryl]x[C6-10 aryl]q[O]p-[C6-10 aryl]-[O]r-[C1-6 alkyl]o-[C(O)]s-[N(R)]-[C(O)]t-[C1-6 alkyl]w- (wherein q, p, r, o, s, t, w, x = 0,1; R = H, C1-4

alkyl, Ph, benzyl), and the remainder of R1, R2, R3, and R4 are individually H, halo, cyano, NO₂, NH₂, sulfonamido, C1-6 alkyl, C1-6 alkoxy, C3-6 cycloalkyl, C3-6 cycloalkyl-C1-6 alkyl, C6-10 aryl, C6-10 aryl-C2-10 alkyl, C6-10 aryl-C2-10 alkenyl, C6-10 heteroaryl, C3-6 heterocycloalkyl, C3-6 heterocycloalkyl-C1-6 alkyl, C2-6 alkenyl, C2-6 alkynyl, C1-6 alkanoyl, halo-C1-6 alkyl, hydroxy-C1-6 alkyl, C1-6 alkoxy, C1-6 alkanoyl, halo-C1-6 alkyl, hydroxy-C1-6 alkyl, C1-6 alkanoyloxy, NR6R7, or SO₂NR6R7; wherein R6, R7 = H, :O, , C1-6 alkyl, C3-6 cycloalkyl, C3-6 cycloalkyl-C1-6 alkyl, Ph, or benzyl; or NR6R7 together form a 5- or 6-membered ring which may optionally contain 1-2 S, (un)substituted NH or nonperoxide O; or R1 and R2 together are methylenedioxy, and optionally any of R1, R2, R3, and R4 is substituted with one to four R1] or pharmaceutically acceptable salts thereof were prepared These compds. I-IV comprise (a) an organic substituent and at least one zinc binding group (ZBG) covalently attached thereto or (b) a ZBG substituted by a side chain. They are inhibitors of metalloprotein, in particular matrix metalloproteinase, histone deacetylase, or anthrax lethal factor, and useful for preventing or treating a pathol. disease, condition, or symptom, in particular cancer, inflammation, or myocardial infarction, that is associated with pathol. metalloprotein activity and/or that is alleviated by inhibition of said activity. Thus, coupling of 3-benzyloxy-6-methyl-4-oxo-4H-pyran-2-carboxylic acid N-(4-iodobenzylamide) with 4-cyanophenylboronic acid in a mixture of aqueous 2 M

aqueous

K₂CO₃ solution and toluene in the presence of Pd(C₂H₃O₂)₂ and PPh₃ under refluxing for 10 days gave 3-benzyloxy-6-methyl-4-oxo-4H-pyran-2-carboxylic acid N-[4-(4-cyanophenyl)benzyl]amide which underwent hydrogenolysis over 10% Pd/C in methanol at 35 psi for 20 h to give 3-hydroxy-6-methyl-4-oxo-4H-pyran-2-carboxylic acid N-[4-(4-cyanophenyl)benzyl]amide (V). V showed IC₅₀ of >50, 0.61, and 0.010 μM against matrix metalloproteinase-1 (MMP-1), MMP-2, and MMP-3, resp.


IT 937187-86-5P, N-[(Biphenyl-4-yl)methyl]-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxamide

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of hydroxydihydropyridinones, hydroxydihydropyridinethiones, hydroxypyranones, and hydroxypyranthiones as metalloprotein inhibitors)

RN 937187-86-5 CAPLUS

CN 2-Pyridinecarboxamide, N-([1,1'-biphenyl]-4-ylmethyl)-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)



L4 ANSWER 3 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:1031424 CAPLUS

DOCUMENT NUMBER: 145:397783

TITLE: Preparation of 3-hydroxypyridin-4-ones as iron modulators

INVENTOR(S): Hider, Robert Charles; Gaeta, Alessandra; Liu, Zu Dong

PATENT ASSIGNEE(S): BTG International Limited, UK

SOURCE: PCT Int. Appl., 66pp.

CODEN: PIXXD2

10/580,011

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006103463	A1	20061005	WO 2006-GB1199	20060331
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

PRIORITY APPLN. INFO.:



GB 2005-6677

A 20050401

OTHER SOURCE(S):

MARPAT 145:397783

GI



AB Title compds. represented by the formula I [wherein R1 = H, (hydroxy)alkyl or (hydroxy)alkenyl; R2 = H, (hydroxy)alkyl, (hydroxy)alkenyl or (un)substituted aralkyl; R3 = H, alkyl, alkenyl or acyl; R4 = H or alkyl; R5-R7 = independently H, (un)substituted alkyl, aryl or aralkyl; R6R7 = (hydroxy)heterocyclyl; and pharmaceutically acceptable tautomers, esters or addition salts thereof] were prepared as iron modulators. For example, II was provided in a multi-step synthesis starting from maltol. I showed relative inhibition of tyrosine hydroxylase, lipoxigenase, and etc.

IT 911289-24-2P 911289-25-3P 911289-26-4P

911289-27-5P 911289-28-6P 911289-29-7P

911289-30-0P 911289-31-1P 911289-32-2P

911289-33-3P 911289-34-4P 911289-36-6P


RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of 3-hydroxypyridin-4-ones as iron modulators)

10/580,011

RN 911289-24-2 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-[2-(methylamino)-2-oxoethyl]-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 911289-25-3 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-[2-[(2-methylpropyl)amino]-2-oxoethyl]-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 911289-26-4 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-[(1S)-1-methyl-2-(methylamino)-2-oxoethyl]-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.




● HCl

RN 911289-27-5 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-[(1S)-1-methyl-2-[(2-methylpropyl)amino]-2-oxoethyl]-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/580,011



● HCl

RN 911289-34-4 CAPLUS


CN 2-Pyridinecarboxamide, N-[2-(dimethylamino)-2-oxoethyl]-1,4-dihydro-3-hydroxy-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 911289-36-6 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-4-oxo-N-[2-oxo-2-(1-piperidinyl)ethyl]-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

REFERENCE COUNT:

2

THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 4 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:1022128 CAPLUS

DOCUMENT NUMBER: 145:356661

TITLE: Preparation of cycloalkyl derivatives of 3-hydroxy-4-pyridinones as therapeutic iron chelating agents

INVENTOR(S): Tam, Tim Fat; Spino, Michael; Li, Wanren; Wang, Yingsheng; Zhao, Yanqing; Shah, Birenkumar Hasmukhbhai

PATENT ASSIGNEE(S): Apotex Inc., Can.

SOURCE: PCT Int. Appl., 77pp.

CODEN: PIXXD2


DOCUMENT TYPE:

Patent

10/580,011

LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005049609	A1	20050602	WO 2004-CA1986	20041118
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
NZ 529657	A	20031219	NZ 2003-529657	20031120
AU 2004291184	A1	20050602	AU 2004-291184	20041118
CA 2546781	A1	20050602	CA 2004-2546781	20041118
EP 1687298	A1	20060809	EP 2004-818733	20041118
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR, IS, YU				
BR 2004016703	A	20070116	BR 2004-16703	20041118
CN 1926135	A	20070307	CN 2004-80039267	20041118
IN 2006MN00572	A	20070309	IN 2006-MN572	20060516
MX 2006PA05594	A	20061219	MX 2006-PA5594	20060517
NO 2006002262	A	20060811	NO 2006-2262	20060519
US 2007082904	A1	20070412	US 2006-580011	20060519
PRIORITY APPLN. INFO.:			NZ 2003-529657	A 20031120
			WO 2004-CA1986	W 20041118
OTHER SOURCE(S):		MARPAT 145:356661		
GI				




AB The invention relates to novel 3-hydroxy-4-pyridinone derivs. of formula I (wherein R1 = X with the proviso that R2 = Y; or R1 = T with the proviso that R2 = W; or R1 = X with the proviso that R2R5N together form an (un)substituted heterocyclic ring; X = C3-C6 cycloalkyl; Y = C3-C6 cycloalkyl, (un)substituted C1-C6 alkyl; T = C1-C6 alkyl; W = C3-C6 cycloalkyl; R3, R4, and R5 = H or C1-C6 alkyl) and their use in chelating ferric (III) ions. Pharmaceutical compns. of such compds. are useful in the removal of excess body iron from patients with iron overload diseases. A process for preparing I is addnl. claimed. For example, II was prepared by hydrogenation of 3-benzyloxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid cyclopropylamide. II at a dose of 450 µmoles/kg in iron overloaded rats caused fecal iron excretion 3 days after administration of 4411 µg/day/kg compared with a baseline value of 3057


10/580,011

ug/day/kg.

- IT 887774-94-9P, 1-Cyclopropyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid methylamide 887774-95-0P, N-(Cyclohexylmethyl)-1-cyclopropyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxamide 887774-96-1P, 1-Cyclopropyl-3-hydroxy-6-methyl-N-(3-methylbutyl)-4-oxo-1,4-dihydropyridine-2-carboxamide 887774-97-2P, 1-Cyclopropyl-N-hexyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxamide 887774-98-3P, N-Cyclohexyl-1-cyclopropyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxamide 887774-99-4P, 1-Cyclopropyl-3-hydroxy-N,N-dimethyl-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxamide 887775-01-1P, 1-Cyclopropyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid cyclopropylamide 910293-45-7P, 3-Hydroxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid cyclohexylamide 910293-46-8P, 3-Hydroxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid cyclopropylamide 910293-51-5P, N-Cyclobutyl-3-hydroxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxamide 910293-52-6P, N-Cyclopentyl-3-hydroxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxamide
- RL: PAC (Pharmacological activity); PRP (Properties); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
- (drug candidate; preparation of cycloalkyl derivs. of 3-hydroxy-4-pyridinones as therapeutic iron chelating agents)
- RN 887774-94-9 CAPLUS
- CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-N,6-dimethyl-4-oxo- (CA INDEX NAME)




- RN 887774-95-0 CAPLUS
- CN 2-Pyridinecarboxamide, N-(cyclohexylmethyl)-1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)




- RN 887774-96-1 CAPLUS
- CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-N-(3-methylbutyl)-4-oxo- (CA INDEX NAME)

10/580,011




RN 887774-97-2 CAPLUS

CN 2-Pyridinecarboxamide, 1-cyclopropyl-N-hexyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)




RN 887774-98-3 CAPLUS

CN 2-Pyridinecarboxamide, N-cyclohexyl-1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)



RN 887774-99-4 CAPLUS


CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-N,N,6-trimethyl-4-oxo- (CA INDEX NAME)




RN 887775-01-1 CAPLUS

CN 2-Pyridinecarboxamide, N,1-dicyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)


10/580,011




RN 910293-45-7 CAPLUS
CN 2-Pyridinecarboxamide, N-cyclohexyl-1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo- (CA INDEX NAME)




RN 910293-46-8 CAPLUS
CN 2-Pyridinecarboxamide, N-cyclopropyl-1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo- (CA INDEX NAME)



RN 910293-51-5 CAPLUS
CN 2-Pyridinecarboxamide, N-cyclobutyl-1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo- (CA INDEX NAME)



RN 910293-52-6 CAPLUS
CN 2-Pyridinecarboxamide, N-cyclopentyl-1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 5 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:1004384 CAPLUS

DOCUMENT NUMBER: 146:14880

TITLE: Fenton Chemistry and Iron Chelation under Physiologically Relevant Conditions: Electrochemistry and Kinetics

AUTHOR(S): Merkofer, Martin; Kissner, Reinhard; Hider, Robert C.; Brunk, Ulf T.; Koppenol, Willem H.

CORPORATE SOURCE: Laboratorium fuer Anorganische Chemie, Departement Chemie und Angewandte Biowissenschaften, ETH Zurich, Zurich, CH-8093, Switz.

SOURCE: Chemical Research in Toxicology (2006), 19(10), 1263-1269

CODEN: CRTOEC; ISSN: 0893-228X

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The goal of Fe-chelation therapy is to reduce the levels of labile plasma Fe, and i.v. administered desferrioxamine is the Au standard of therapeutic agents. Hydroxypyridinones, e.g., CP20 (3-hydroxy-1,2-dimethylpyridin-4(1H)-one), are used or are under study as orally administered Fe chelators. The authors determined electrode potentials of CP20, the related hydroxypyridones CP361, CP363, and CP502, and ICL670 (4-[3,5-bis(2-hydroxyphenyl)-1H-1,2,4-triazol-1-yl]benzoic acid) under physiol. relevant conditions to address the question of whether Fe in the presence of these chelating agents can carry out Fenton chemical in vivo. Fe(III) but not Fe(II) binds tightly to both CP20 and ICL670 at pH 7 and higher, compared to nearly complete binding of 1 μ M Fe(II) to 10 μ M desferrioxamine at pH 7.4. The electrode potentials of the hydroxypyridinones shift to more neg. values with decreasing pKa values at lower concns. of Fe(III) (0.02 mM) and ligand (0.1 mM). The electrode potential of the Fe-CP20 system decreases as a function of increasing pH, with a min. near pH 10.5. The authors estimate an electrode potential for the ascorbyl radical/ascorbate couple under physiol. conditions of +105 mV, which is higher than the electrode potential of the Fe(III) complex of CP20 at all concns. of Fe. The rate of oxidation of Fe(II) in the presence of CP20 by H₂O₂ increases with the concns. of both ligand and peroxide. Although Fe(II) is oxidized by H₂O₂, the thus-formed FeIII(CP20)₃ complex cannot be reduced by ascorbate. Therefore, the tight binding of Fe(III) by this class of chelators prevents redox cycling.

IT 243987-44-2D, CP 502, iron complexes


RL: CPS (Chemical process); PEP (Physical, engineering or chemical process); PRP (Properties)

(cyclic voltammetry on mercury drop electrode in Tris buffer and reaction kinetics with H₂O₂ and Fenton reaction and iron chelation under physiol. relevant conditions and electrochem. and kinetics)


RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)

10/580,011



IT 793651-87-3, CP 509
RL: PRP (Properties)
(reaction kinetics with H2O2 and Fenton reaction and iron chelation
under physiol. relevant conditions and electrochem. and kinetics)
RN 793651-87-3 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,N,1,6-tetramethyl-4-oxo-
(CA INDEX NAME)




REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 6 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2006:485216 CAPLUS
DOCUMENT NUMBER: 145:8031
TITLE: Process for the preparation of 3-hydroxy-1-cycloalkyl-
6-alkyl-4-oxo-1,4-dihydropyridine-2-carboxamides by
treatment of the corresponding acids with acid
chloride formation reagents and amines.
INVENTOR(S): Wang, Yingsheng; Agostino, Sandra Vittoria; Tam, Tim
Fat; Zhao, Yanqing; Li, Wanren; Shah, Birenkumar
Hasmukhbhai; Leung-Toung, Regis
PATENT ASSIGNEE(S): Apotex Inc., Can.
SOURCE: Can. Pat. Appl., 45 pp.
CODEN: CPXXEB
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2488034	A1	20060519	CA 2004-2488034	20041119
WO 2006053429	A1	20060526	WO 2005-CA1746	20051117
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				

IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ,
 CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,
 GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,
 KG, KZ, MD, RU, TJ, TM
 EP 1824823 A1 20070829 EP 2005-810777 20051117
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,
 IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR
 CN 101094834 A 20071226 CN 2005-80045616 20051117
 IN 2007DN04443 A 20070831 IN 2007-DN4443 20070611
 PRIORITY APPLN. INFO.: CA 2004-2488034 A 20041119
 WO 2005-CA1746 W 20051117
 OTHER SOURCE(S): CASREACT 145:8031; MARPAT 145:8031
 GI




AB Title compds. [I; R1, R4 = H, alkyl; R2 = alkyl, cycloalkyl; R3 = alkyl, cycloalkyl, H, cycloalkylalkyl; R5 = H, (substituted) PhCH2, protecting group; R6 = H, alkyl, cycloalkyl; NR3R6 = (substituted) piperidinyl, morpholinyl, pyrrolidinyl, piperazinyl], were prepared via reaction of the corresponding acids with acid chloride formation reagents and amines. Thus, 3-benzyloxy-1-cyclopropyl-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid (preparation given) and DMF in CH2Cl2 were treated with (COCl)2 over 1 h at <10°. The resulting solution was added over 2.5 h to a solution of Et3N and MeNH2 in THF at 4° to give 90% 3-benzyloxy-1-cyclopropyl-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid methylamide. The latter was hydrogenated in MeOH/H2O containing concentrate HCl over Pd/C under 50 psi H2 for 3 h to give 74% 1-cyclopropyl-3-hydroxy-6-methyl-4-oxo-1,4-dihydropyridine-2-carboxylic acid methylamide.

IT 887774-94-9P 887774-95-0P 887774-96-1P
 887774-97-2P 887774-98-3P 887774-99-4P
 887775-01-1P
 RL: IMF (Industrial manufacture); SPN (Synthetic preparation); PREP (Preparation)
 (preparation of hydroxycycloalkylalkyloxodihydropyridinecarboxamides by treatment of the corresponding acids with acid chloride formation reagents and amines)


RN 887774-94-9 CAPLUS
 CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-N,6-dimethyl-4-oxo- (CA INDEX NAME)

10/580,011




RN 887774-95-0 CAPLUS

CN 2-Pyridinecarboxamide, N-(cyclohexylmethyl)-1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)




RN 887774-96-1 CAPLUS

CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-N-(3-methylbutyl)-4-oxo- (CA INDEX NAME)



RN 887774-97-2 CAPLUS


CN 2-Pyridinecarboxamide, 1-cyclopropyl-N-hexyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)



RN 887774-98-3 CAPLUS


CN 2-Pyridinecarboxamide, N-cyclohexyl-1-cyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)

10/580,011




RN 887774-99-4 CAPLUS

CN 2-Pyridinecarboxamide, 1-cyclopropyl-1,4-dihydro-3-hydroxy-N,N,6-trimethyl-4-oxo- (CA INDEX NAME)



RN 887775-01-1 CAPLUS

CN 2-Pyridinecarboxamide, N,1-dicyclopropyl-1,4-dihydro-3-hydroxy-6-methyl-4-oxo- (CA INDEX NAME)



L4 ANSWER 7 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:453925 CAPLUS

DOCUMENT NUMBER: 145:76021

TITLE: Structure-activity relationship studies on UK-2A, a novel antifungal antibiotic from *Streptomyces* sp. 517-02. Part 5: Roles of the 9-membered dilactone-ring moiety in respiratory inhibition

AUTHOR(S): Usuki, Yoshinosuke; Adachi, Noriko; Fujita, Ken-Ichi; Ichimura, Akio; Iio, Hideo; Taniguchi, Makoto

CORPORATE SOURCE: Department of Material Science, Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi, Osaka, 558-8585, Japan

SOURCE: Bioorganic & Medicinal Chemistry Letters (2006), 16(12), 3319-3322

CODEN: BMCLE8; ISSN: 0960-894X

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 145:76021

AB Several open-chained analogs of UK-2A, a novel antifungal antibiotic isolated from *Streptomyces* sp. 517-02, were prepared for structure-activity

10/580,011

studies. The in vitro antifungal activities of these compds. against *Rhodotorula mucilaginosa* IFO 0001 and the inhibition of uncoupler-stimulated respiration in bovine heart submitochondrial particles (SMP) were evaluated. Oxidative potentials were measured by cyclic voltammetry. An analog prepared from dihexyl -glutamate showed comparable inhibitory activity as UK-2A.

IT 894354-57-5P 894354-58-6P 894354-59-7P

894354-60-0P 894354-61-1P 894354-62-2P


RL: DMA (Drug mechanism of action); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(SAR of and preparation antifungal UK-2A analogs: 9-membered dilactone-ring moiety role in respiratory inhibition)

RN 894354-57-5 CAPLUS

CN L-Glutamic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, dipentyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 894354-58-6 CAPLUS

CN L-Glutamic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, dihexyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 894354-59-7 CAPLUS

CN L-Glutamic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, diheptyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.


10/580,011



RN 894354-60-0 CAPLUS

CN L-Glutamic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, dioctyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 894354-61-1 CAPLUS

CN D-Glutamic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, dihexyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 894354-62-2 CAPLUS

CN L-Aspartic acid, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, dihexyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 8 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:331313 CAPLUS

DOCUMENT NUMBER: 145:3116

TITLE: Discovery of Different Types of Inhibition between the Human and *Thermotoga maritima* α -Fucosidases by Fuconojirimycin-Based Derivatives

AUTHOR(S): Ho, Ching-Wen; Lin, Yu-Nong; Chang, Chuan-Fa; Li, Shiou-Ting; Wu, Ying-Ta; Wu, Chung-Yi; Chang, Chiung-Fang; Liu, Sheng-Wen; Li, Yaw-Kuen; Lin, Chun-Hung

CORPORATE SOURCE: Institute of Biological Chemistry and Genomics Research Center, Academia Sinica, Taipei, 11529, Taiwan

SOURCE: Biochemistry (2006), 45(18), 5695-5702

CODEN: BICHAW; ISSN: 0006-2960

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An efficient method for examining the selectivity of inhibitors on two α -fucosidases, one from *Thermotoga maritima* and the other from human, was established. The X-ray crystal structure of the former enzyme makes possible the homol. modeling of the human α -fucosidase, indicating the major difference between both enzymes in the periphery of the catalytic site. To investigate the difference at the mol. level, a variety of fuconojirimycin (FNJ) derivs. with substitution at C1, C2, C6, or N were rapidly prepared in microplates and screened without purification for the inhibition activities of the two α -fucosidases. Among the mols. that were tested, only the substitution at C1 can significantly enhance the inhibitory potency, in contrast to the control (no substitution) and compds. with substitution at other positions. The majority of C1-substituted FNJs were found to be slow tight-binding inhibitors of the *Thermotoga* enzyme, while acting as the reversible inhibitors of the human fucosidase. The best inhibitor exhibited 13,700-fold difference in affinity between the two enzymes, which was attributed to the dissimilar aglycon binding site. Further investigations were carried out, including site-directed mutagenesis, the comparison of K_i values among the wild-type and mutants, and the intrinsic fluorescence change upon inhibitor titration, all supporting the idea that Tyr64 and Tyr267 of the *Thermotoga* α -fucosidase are critically involved in closely interacting with the aglycon of inhibitors. The increased level of contact thus induced conformational change, leading to the observed slow tight-binding inhibition.

IT 887948-57-4P 887949-34-0P 887950-11-0P

RL: BSU (Biological study, unclassified); PRP (Properties); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)


(preparation of fuconojirimycin derivs. as inhibitors; different types of inhibition between the human and *Thermotoga maritima* α -fucosidases by fuconojirimycin-based derivs.)

10/580,011

RN 887948-57-4 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-methyl-N-[[[(2R,3R,4R,5R,6S)-3,4,5-trihydroxy-6-methyl-2-piperidinyl]methyl]-, (2S,3R,4R,5R,6S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 887949-34-0 CAPLUS

CN 2-Piperidinecarboxamide, N-[3-[[[(3R,4R,5R,6S)-4,5-dihydroxy-6-methyl-3-piperidinyl]oxy]propyl]-3,4,5-trihydroxy-6-methyl-, (2S,3R,4R,5R,6S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 887950-11-0 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-methyl-N-[3-[(2S,3R,4S,5R)-3,4,5-trihydroxy-2-methyl-1-piperidinyl]propyl]-, (2S,3R,4R,5R,6S)- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

44

THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L4 ANSWER 9 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:272910 CAPLUS

DOCUMENT NUMBER: 144:331269

TITLE: Preparation of carbamoylpyridone derivative having HIV integrase inhibitory activity
 INVENTOR(S): Yoshida, Hiroshi
 PATENT ASSIGNEE(S): Shionogi & Co., Ltd., Japan
 SOURCE: PCT Int. Appl., 87 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006030807	A1	20060323	WO 2005-JP16904	20050914
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
EP 1790638	A1	20070530	EP 2005-783196	20050914
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR			
CN 101014572	A	20070808	CN 2005-80030458	20050914
US 2007249687	A1	20071025	US 2007-662768	20070314
IN 2007CN01084	A	20070907	IN 2007-CN1084	20070315
PRIORITY APPLN. INFO.:			JP 2004-267720	A 20040915
			WO 2005-JP16904	W 20050914
OTHER SOURCE(S):		MARPAT 144:331269		
GI				



AB Title compds. represented by the formula I [wherein Y = (un)substituted N, O, S or SO₂; R = COR₅ or (un)substituted N containing cyclyl; R₂, R₅ = independently H, OH, alkyl, etc.; X = single bond, O, S, SO, etc.; R₃ = H, halo, OH, (un)substituted alkyl, etc.; and pharmaceutically acceptable salts or solvates thereof] were prepared as anti-HIV agents. For example, II was provided in a multi-step synthesis starting from 4-hydroxy-6-methylnicotinic acid. II showed inhibition of integrase with IC₅₀ value of 6.4 ng/mL. Thus, I are useful as antiviral agents for the treatment of HIV.

IT 880261-29-0P 880261-30-3P 880261-31-4P

10/580,011


880261-32-5P 880261-33-6P 880261-34-7P
880261-36-9P 880261-37-0P 880261-38-1P
880261-39-2P 880261-40-5P 880261-41-6P
880261-43-8P 880261-56-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU
(Therapeutic use); BIOL (Biological study); PREP (Preparation); USES
(Uses)

(preparation of carbamoylpyridone derivative having HIV integrase inhibitory
activity)


RN 880261-29-0 CAPLUS

CN 2,5-Pyridinedicarboxamide, N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-
hydroxy-N2-methyl-4-oxo- (CA INDEX NAME)




RN 880261-30-3 CAPLUS

CN 2,5-Pyridinedicarboxamide, N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-
hydroxy-N2,N2-dimethyl-4-oxo- (CA INDEX NAME)



RN 880261-31-4 CAPLUS


CN 2,5-Pyridinedicarboxamide, N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-
hydroxy-N2-(1-methylethyl)-4-oxo- (CA INDEX NAME)



RN 880261-32-5 CAPLUS

CN 2,5-Pyridinedicarboxamide, N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-
hydroxy-N2-(2-methoxyethyl)-4-oxo- (CA INDEX NAME)


10/580,011



RN 880261-41-6 CAPLUS


CN 2,5-Pyridinedicarboxamide, N2-[(1S)-1-cyclohexylethyl]-N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-hydroxy-4-oxo- (CA INDEX NAME)

Absolute stereochemistry.




RN 880261-43-8 CAPLUS

CN 2,5-Pyridinedicarboxamide, N2-[2-(1-cyclohexen-1-yl)ethyl]-N5-[(4-fluorophenyl)methyl]-1,4-dihydro-3-hydroxy-4-oxo- (CA INDEX NAME)



RN 880261-56-3 CAPLUS

CN 2,5-Pyridinedicarboxamide, N,N'-bis[(4-fluorophenyl)methyl]-1,4-dihydro-3-hydroxy-4-oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L4 ANSWER 10 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2006:261710 CAPLUS

10/580,011

DOCUMENT NUMBER: 145:28251
TITLE: Preparation of α -ketoamide or hydroxyethylamine
peptidomimetics as HIV protease inhibitors
INVENTOR(S): Laslo, Karen; Slee, Deborah H.; Wong, Chi-Huey
PATENT ASSIGNEE(S): The Scripps Research Institute, Australia
SOURCE: Aust. Pat. Appl., 191 pp., Division of Aust. 2001
18,270.
CODEN: AUXXCM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
AU 2004202175	A1	20040617	AU 2004-202175	20040520
PRIORITY APPLN. INFO.:			AU 2001-18270	A3 20010202
OTHER SOURCE(S):	MARPAT 145:28251			
GI				



AB Combinatorial libraries of HIV and FIV protease inhibitors are characterized by α -ketoamide or hydroxyethylamine core structures flanked by on one side by substituted pyrrolidines, piperidines, or aza sugars and on the other side by phenylalanine, tyrosine, or substituted tyrosines. α -Ketoamide I [R₁ is H, OH, alkoxy, OBn or OP, where P is a protecting group; R₂, R₃ are independently groups given for R₁ or benzyloxy substituted by methoxy, nitro or hydroxy groups; R₄ is H, CH₂OH, alkoxyethyl or CH₂OP (with the proviso that R₁-R₄ cannot all be H)] are claimed. The libraries are synthesized via a one step coupling reaction. Highly efficacious drug candidates are identified by screening the libraries for binding and inhibitory activity against both HIV and FIV protease. Drug candidates displaying clin. useful activity against both HIV and FIV protease are identified as being potentially resistive against a loss of inhibitory activity due to development of resistant strains of

10/580,011

HIV. Thus, α -ketoamide II (Cbz = benzyloxycarbonyl) was prepared and showed $K_i = 65$ nM for inhibition of HIV protease.


IT 191850-39-2 191850-42-7 191850-45-0
191850-48-3

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation of α -ketoamide or hydroxyethylamine peptidomimetics as HIV protease inhibitors)

RN 191850-39-2 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3R,4R,5S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-42-7 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3S,4R,5S)- (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-45-0 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3R,4R,5R)- (CA INDEX NAME)

Absolute stereochemistry.




RN 191850-48-3 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3S,4R,5R)- (CA INDEX NAME)

Absolute stereochemistry.

10/580,011



IT 888948-52-5P


RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of α -ketoamide or hydroxyethylamine peptidomimetics as HIV protease inhibitors)

RN 888948-52-5 CAPLUS

CN 1-Piperidinecarboxylic acid, 2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-, phenylmethyl ester, (2S,3S,4R,5S)- (CA INDEX NAME)

Absolute stereochemistry.



IT 191850-51-8P 191850-64-3P 191850-67-6P

191850-75-6P


RL: SPN (Synthetic preparation); PREP (Preparation)

(preparation of α -ketoamide or hydroxyethylamine peptidomimetics as HIV protease inhibitors)

RN 191850-51-8 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3R,4R,5S)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 191850-64-3 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3S,4R,5S)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

10/580,011 .


Absolute stereochemistry.



RN 191850-67-6 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3R,4R,5R)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-75-6 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3S,4R,5R)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 11 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2006:140501 CAPLUS
DOCUMENT NUMBER: 144:343023

10/580,011

TITLE: Transport kinetics of iron chelators and their chelates in Caco-2 cells

AUTHOR(S): Huang, Xi-Ping; Spino, M.; Thiessen, J. J.

CORPORATE SOURCE: Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON, M5S 2S2, Can.

SOURCE: Pharmaceutical Research (2006), 23(2), 280-290
CODEN: PHREEB; ISSN: 0724-8741

PUBLISHER: Springer

DOCUMENT TYPE: Journal


LANGUAGE: English

AB Caco-2 monolayers were used to contrast the bidirectional transport of iron chelators and their chelates and to estimate fundamental kinetics associated with their intestinal absorption. Bidirectional transport was studied at 37°C and pH 7.4 using 500-µM concns. Monolayer integrity was tested via transepithelial elec. resistance and sodium fluorescein permeability. Apical and basolateral anal. provided mass balance evidence. Apparent permeability coefficient (Papp) served to rank and compare mols. and estimate in vivo bioavailability. Model-dependent rate consts. defined cellular influx and efflux. Papp ranked in decreasing order for chelators from directional transport studies were CP363 > deferiprone > ICL670 > CP502 > deferoxamine (DFO). Fe(CP502)3, Fe(ICL670)2, and FeDFO were not measurable in receiving chambers, whereas Fe(deferiprone)3 and Fe(CP363)3 were detected in both directions. CP363 was transported significantly faster from the basolateral to the apical direction than the converse. Mass balance of donor and receiver chambers gave approx. 100% recovery in all cases. Kinetic anal. supports the view that the Caco-2 chelator efflux consts. are generally greater than their influx consts. Caco-2 cells are useful in screening iron chelators and chelates and estimating bioavailabilities. Structure and distribution coeffs. partially predict passive transport through Caco-2 monolayers.

IT 243987-44-2
RL: PKT (Pharmacokinetics); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(transport kinetics of iron chelators and their chelates in Caco-2 cells)

RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 12 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:1084925 CAPLUS

DOCUMENT NUMBER: 144:23184

TITLE: High affinity iron(III) scavenging by a novel hexadentate 3-hydroxypyridin-4-one-based dendrimer: Synthesis and characterization

AUTHOR(S): Zhou, Tao; Liu, Zu Dong; Neubert, Hendrik; Kong, Xiao Le; Ma, Yong Min; Hider, Robert C.

CORPORATE SOURCE: Department of Pharmacy, King's College London, London,

SE1 9NH, UK
 SOURCE: Bioorganic & Medicinal Chemistry Letters (2005),
 15(22), 5007-5011
 CODEN: BMCLE8; ISSN: 0960-894X
 PUBLISHER: Elsevier B.V.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 144:23184


AB The synthesis of a novel iron(III)-selective hydroxypyridinone hexadentate-terminated dendritic chelator based on a benzene tricarbonyl core polyamine dendrimer is described. The iron-chelating ability of the dendritic chelator was demonstrated by matrix-assisted laser desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS) and UV-vis spectroscopy. The physicochem. properties of the isolated hexadentate unit were also investigated. The dendrimer was found to possess an extremely high affinity for iron(III), namely $\log K = 34.8$, $pFe_{3+} = 30.6$.



IT 870456-70-5P
 RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (dendritic; preparation and characterization of high affinity iron(III) scavenging by a novel hexadentate 3-hydroxypyridin-4-one-based dendrimer)

RN 870456-70-5 CAPLUS


CN 1,3,5-Benzenetricarboxamide, N,N',N''-tris[4-[[[(1,4-dihydro-3-hydroxy-4-oxo-2-pyridinyl)carbonyl]amino]-1,1-bis[3-[[[(1,4-dihydro-3-hydroxy-4-oxo-2-pyridinyl)carbonyl]amino]propyl]butyl]- (9CI) (CA INDEX NAME)

PAGE 1-A





IT 870456-67-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and characterization of high affinity iron(III) scavenging by a
 novel hexadentate 3-hydroxypyridin-4-one-based dendrimer)
 RN 870456-67-0 CAPLUS
 CN 2-Pyridinecarboxamide, N,N'-[4-[3-[[[(1,4-dihydro-3-hydroxy-4-oxo-2-
 pyridinyl) carbonyl] amino] propyl] -4-nitro-1,7-heptanediyl] bis[1,4-dihydro-3-
 hydroxy-4-oxo- (9CI) (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 13 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:823463 CAPLUS

DOCUMENT NUMBER: 143:229725

TITLE: Preparation of N-Benzyl-3,4-dihydroxypyridine-2-carboxamides useful as HIV integrase inhibitors
INVENTOR(S): Jones, Philip; Williams, Peter D.; Morrisette, Matthew M.; Kuo, Michelle Sparks; Vacca, Joseph P.
PATENT ASSIGNEE(S): Merck & Co., Inc., USA; Istituto di Ricerche di Biologia Molecolare P. Angeletti S.p.A.

SOURCE: PCT Int. Appl., 81 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005074513	A2	20050818	WO 2005-US2472	20050126
WO 2005074513	A3	20050929		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2005211349	A1	20050818	AU 2005-211349	20050126
CA 2554120	A1	20050818	CA 2005-2554120	20050126
EP 1713773	A2	20061025	EP 2005-726383	20050126
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK, IS			
JP 2007519735	T	20070719	JP 2006-551441	20050126
CN 101014571	A	20070808	CN 2005-80003386	20050126
US 2007155744	A1	20070705	US 2006-587330	20060727
IN 2006DN04345	A	20070713	IN 2006-DN4345	20060727
PRIORITY APPLN. INFO.:			US 2004-540538P	P 20040130
			WO 2005-US2472	W 20050126

10/580,011

OTHER SOURCE(S): CASREACT 143:229725; MARPAT 143:229725

AB N-Benzyl-dihydroxypyridine carboxamide compds. are inhibitors of HIV integrase and inhibitors of HIV replication. The compds. are useful in the prevention and treatment of infection by HIV and in the prevention, delay in the onset, and treatment of AIDS. The compds. and their salts can be employed as ingredients in pharmaceutical compns., optionally in combination with other antivirals, immunomodulators, antibiotics or vaccines. An example compound prepared was 6-acetyl-N-(4-fluorobenzyl)-3,4-dihydroxypyridine-2-carboxamide.

IT 862667-37-6P 862667-38-7P 862667-39-8P


862667-40-1P 862667-41-2P 862667-44-5P

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation of N-benzyl-3,4-dihydroxypyridine-2-carboxamides useful as HIV integrase inhibitors)


RN 862667-37-6 CAPLUS

CN 2-Pyridinecarboxamide, 6-acetyl-N-[(4-fluorophenyl)methyl]-3,4-dihydroxy-
(CA INDEX NAME)




RN 862667-38-7 CAPLUS

CN 2-Pyridinecarboxylic acid, 6-[[[(4-fluorophenyl)methyl]amino]carbonyl]-4,5-dihydroxy- (CA INDEX NAME)




RN 862667-39-8 CAPLUS

CN 2-Pyridinecarboxylic acid, 6-[[[(4-fluorophenyl)methyl]amino]carbonyl]-4,5-dihydroxy-, methyl ester (CA INDEX NAME)



RN 862667-40-1 CAPLUS

10/580,011



L4 ANSWER 14 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:409512 CAPLUS

DOCUMENT NUMBER: 142:463613

TITLE: A preparation of pyridinecarboxamide derivatives, useful for inhibiting HIV integrase

INVENTOR(S): Kong, Laval Chan Chun; Zhang, Ming-Qiang; Halab, Liliane; Nguyen-Ba, Nghe; Liu, Bingcan

PATENT ASSIGNEE(S): Virochem Pharma Inc., Can.

SOURCE: PCT Int. Appl., 139 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:



PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005042524	A1	20050512	WO 2004-CA1898	20041029
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				

US 2005176767 A1 20050811 US 2004-976238 20041029

PRIORITY APPLN. INFO.: US 2003-515443P P 20031030

OTHER SOURCE(S): CASREACT 142:463613; MARPAT 142:463613

GI




AB The invention relates to a preparation of pyridinecarboxamide derivs. of formula I [wherein: R1 is H or alkyl; R2 is OH, alkoxy, or arylalkoxy; R3 is NH2, amido, sulfonamido, azido, halogen, or alkoxy, etc.; R4 is H, halogen, OH, carboxy, or (hetero)aryl, etc.; R5 and R6 are independently H, alkyl, aryl, or arylalkyl; Q is (un)substituted Ph, alkyl, or heterocyclyl, etc.], useful for inhibiting HIV integrase. For instance, pyridinecarboxamide derivative II was prepared via Pd-catalyzed coupling of the prepared 4-iodo-3-hydroxypyridine-2-carboxylic acid 4-fluorobenzylamide with 2-trimethylstannyl-pyridine. Certain invention compds. were tested in an assay for HIV activity (IC50 <10 μ M).

IT 851441-89-9P, 6-Bromo-3,4-dihydroxypyridine-2-carboxylic acid 4-fluorobenzylamide 851442-00-7P, 6-Furan-2-yl-3-hydroxy-4-methoxypyridine-2-carboxylic acid 4-fluorobenzylamide 851442-63-2P
 RL: PAC (Pharmacological activity); RCT (Reactant); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
 (preparation of pyridinecarboxamide derivs. useful for inhibiting HIV integrase)

RN 851441-89-9 CAPLUS


CN 2-Pyridinecarboxamide, 6-bromo-N-[(4-fluorophenyl)methyl]-3,4-dihydroxy-
 (CA INDEX NAME)



RN 851442-00-7 CAPLUS


CN 2-Pyridinecarboxamide, N-[(4-fluorophenyl)methyl]-6-(2-furanyl)-3-hydroxy-4-methoxy- (CA INDEX NAME)

10/580,011



RN 851443-00-0 CAPLUS

CN 2-Pyridinecarboxamide, N-[(4-fluorophenyl)methyl]-3-hydroxy-4-methoxy-6-(5-methyl-1,3-oxathian-2-yl)- (CA INDEX NAME)




IT 851442-14-3, 4-Benzyloxy-3-hydroxy-6-vinylpyridine-2-carboxylic acid 4-fluorobenzylamide

RL: RCT (Reactant); RACT (Reactant or reagent)

(reactant; preparation of pyridinecarboxamide derivs. useful for inhibiting HIV integrase)

RN 851442-14-3 CAPLUS

CN 2-Pyridinecarboxamide, 6-ethenyl-N-[(4-fluorophenyl)methyl]-3-hydroxy-4-(phenylmethoxy)- (CA INDEX NAME)



REFERENCE COUNT:

5

THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 15 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:158668 CAPLUS

DOCUMENT NUMBER: 142:261561



TITLE: Preparation of pyrido[1,2-a]pyrazine-1,8-dione derivatives as HIV integrase inhibitors

INVENTOR(S): Miyazaki, Susumu; Katoh, Susumu; Adachi, Kaoru; Isoshima, Hiroataka; Kobayashi, Satoru; Matsuzaki, Yuji; Watanabe, Wataru; Yamataka, Kazunobu; Kiyonari,

Shinichi; Wamaki, Shuichi
 PATENT ASSIGNEE(S): Japan Tobacco Inc., Japan
 SOURCE: PCT Int. Appl., 355 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005016927	A1	20050224	WO 2004-JP11869	20040812
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2577239	A1	20050224	CA 2004-2577239	20040812
EP 1544199	A1	20050622	EP 2004-771830	20040812
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR				
JP 3814631	B2	20060830	JP 2005-513209	20040812
US 2005054645	A1	20050310	US 2004-958225	20041005
US 2006052361	A1	20060309	US 2005-255605	20051013
US 7211572	B2	20070501		
JP 2006232849	A	20060907	JP 2006-118260	20060421
PRIORITY APPLN. INFO.:				
			JP 2003-293117	A 20030813
			JP 2004-134896	A 20040428
			JP 2005-513209	A3 20040812
			WO 2004-JP11869	W 20040812
			US 2004-958225	B1 20041005

OTHER SOURCE(S): MARPAT 142:261561
 GI




AB The title compds. I [wherein R1 = (un)substituted alkyl, alkenyl, alkynyl, etc.; X = (un)substituted CH2, N=CH, or CH=N; Y1-Y2-Y3 = (un)substituted C=CH-NH, N-CH=N, N-CH=CH, C=N-NH, N-N=CH, etc.; R2 = H, alkyl, arylalkyl, or (un)substituted SO2H] or pharmaceutically acceptable salts thereof are prepared as anti-HIV agents. For example, the compound II•HCl was prepared in a multi-step synthesis. II•HCl inhibited HIV integrase with IC50 of <0.01 μM. Formulations containing I as an active ingredient were also described.

IT 845723-64-0P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (intermediate; preparation of pyrido[1,2-a]pyrazine-1,8-dione derivs. as HIV integrase inhibitors)

10/580,011

RN 845723-64-0 CAPLUS

CN 2-Pyridinecarboxamide, N-[(3-chlorophenyl)methyl]-1-(2,2-dimethoxyethyl)-1,4-dihydro-3-hydroxy-5-(1-methylethyl)-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 16 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:146262 CAPLUS

DOCUMENT NUMBER: 142:329432

TITLE: Metabolic and pharmacokinetic evaluation of a novel 3-hydroxypyridinone iron chelator, CP502, in the rat

AUTHOR(S): Novakovic, Jasmina; Tesoro, Angelo; Thiessen, Jake J.; Spino, Michael

CORPORATE SOURCE: Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON, Can.

SOURCE: European Journal of Drug Metabolism and Pharmacokinetics (2004), 29(4), 221-224
CODEN: EJDPD2; ISSN: 0378-7966

PUBLISHER: Medecine et Hygiene

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A recently synthesized 3-hydroxypyridinone derivative with an amido function at the 2-position, CP502 (1,6-dimethyl-3-hydroxy-4-(1H)-pyridinone-2-carboxy-(N-methyl)-amide hydrochloride), exhibited high in vitro iron chelating potency ($pFe_{3+} = 21.7$). It was targeted as a new iron-chelating candidate for further development in early pre-clin. testing. To evaluate its pharmacokinetics, including oral bioavailability, metabolic and disappearance profiles, studies were conducted in Sprague Dawley male rats. A single 150 mg/kg i.v. and oral dose was given to male Sprague Dawley rats (N=6, B.Weight 250g). The rats were placed in metabolic cages and fasted overnight before the dosing. Venous blood samples (200 μ L per withdrawal) were collected at defined time points before (blank) and up to 28 h post administration. Urine and feces were collected before dosing (blank) and in 24 h intervals up to 72 h post administration. Plasma CP502 concentration vs. time profiles were consistent with 2-compartment distribution, and the oral bioavailability approached 100%. Total clearance and mean residence time (i.v.) were 1.02 L/kg/h and 1.10 h, resp. Simultaneous computer fitting yielded V_1 and V_{ss} ests. of 0.96 L/kg and 1.74 L/kg, resp. CP502 was mainly excreted unchanged via urine (45.29 \pm 9.40 % of total dose) or as glucuronide (6.46 \pm 1.22% of total dose). High iron chelation potential and favorable pharmacokinetic and metabolic profiles indicate that CP502 is a promising candidate for further development.


IT 243987-44-2

RL: PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(metabolic and pharmacokinetic evaluation of 3-hydroxypyridinone iron chelator CP502 in rats)

RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 17 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:85982 CAPLUS

DOCUMENT NUMBER: 142:336222

TITLE: Design and characterization of novel hexadentate 3-hydroxypyridin-4-one ligands

AUTHOR(S): Piyamongkol, Sirivipa; Zhou, Tao; Liu, Zu D.; Khodr, Hicham H.; Hider, Robert C.

CORPORATE SOURCE: Department of Pharmacy, King's College London, London, SE1 9NN, UK

SOURCE: Tetrahedron Letters (2005), 46(8), 1333-1336

CODEN: TELEAY; ISSN: 0040-4039


PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 142:336222

GI



AB Two novel hexadentate 3-hydroxy-4-pyridinone ligands have been designed and synthesized. The physico-chemical properties of one of the hexadentate ligands have been determined and the results indicate that the hexadentate ligand possesses high affinity for iron(III). One of the target compds. prepared for this study was N,N',N''-(nitritotri-2,1-ethanediyl)tris[3,4-di(hydroxy)-2-pyridinecarboxamide] tetrahydrochloride (I). The stability constant of I-iron complex was determined. The applicability of I as potential therapeutic iron chelator is under investigation (no data).


10/580,011

IT 848644-97-3P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
(preparation of N,N',N''-(nitrilotri-2,1-ethanediyl)tris[di(hydroxy)-2-pyridinecarboxamide], study of its iron complex and its stability constant, and its applicability as potential therapeutic iron chelator)

RN 848644-97-3 CAPLUS

CN 2-Pyridinecarboxamide, N,N',N''-(nitrilotri-2,1-ethanediyl)tris[3,4-dihydroxy-, tetrahydrochloride (9CI) (CA INDEX NAME)



●4 HCl


IT 848644-96-2P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of

N,N',N''-[1,3,5-benzenetriyltris(methylene)]tris[di(hydroxy)-2-pyridinecarboxamide] and study of its applicability as ligand for iron)

RN 848644-96-2 CAPLUS

CN 2-Pyridinecarboxamide, N,N',N''-[1,3,5-benzenetriyltris(methylene)]tris[3,4-dihydroxy-, trihydrochloride (9CI) (CA INDEX NAME)




● 3 HCl

IT 349141-36-2P

RL: PRP (Properties); SPN (Synthetic preparation); PREP (Preparation)
 (preparation of di(hydroxy)-N-[(hydroxy)ethyl]pyridinecarboxamide, study of
 its iron complex and determination of its stability constant)

RN 349141-36-2 CAPLUS

CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-(2-hydroxyethyl)-,
 monohydrochloride (9CI) (CA INDEX NAME)



● HCl

REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 18 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2005:39639 CAPLUS

DOCUMENT NUMBER: 142:273946

TITLE: Redox properties of the iron complexes of orally
 active iron chelators CP20, CP502, CP509, and ICL670

AUTHOR(S): Merkofer, Martin; Kissner, Reinhard; Hider, Robert C.;
 Koppenol, Willem H.

CORPORATE SOURCE: Laboratorium fuer Anorganische Chemie, Department
 Chemie und Angewandte Biowissenschaften, Zurich,
 CH-8093, Switz.


SOURCE: Helvetica Chimica Acta (2004), 87(12), 3021-3034
 CODEN: HCACAV; ISSN: 0018-019X

PUBLISHER: Verlag Helvetica Chimica Acta


DOCUMENT TYPE: Journal

LANGUAGE: English

- AB Redox cycling of iron is a critical aspect of iron toxicity. Reduction of a low-mol.-weight iron(III)-complex followed by oxidation of the iron(II)-complex by hydrogen peroxide may yield the reactive hydroxyl radical (OH) or an oxo iron(IV) species (the Fenton reaction). Complexation of iron by a ligand that shifts the electrode potential of the complex to either to far below -350 mV (dioxygen/superoxide, pH = 7) or to far above +320 mV (H₂O₂/HO, H₂O pH = 7) is essential for limiting Fenton reactivity. The oral chelating agents CP20, CP502, CP509, and ICL670 effectively remove iron from patients suffering from iron overload. We measured the electrode potentials of the iron(III) complexes of these drugs by cyclic voltammetry with a mercury electrode and determined the dependence on concentration, pH, and stoichiometry. The standard electrode potentials measured are -620 mV, -620 mV, -535 mV, and -535 mV with iron bound to CP20, ICL670, CP502, and CP509, resp., but, at lower chelator concns., electrode potentials are significantly higher.
- IT 243987-44-2D, iron complexes 793651-87-3D, iron complexes
 RL: BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)
 (redox properties of iron complexes of orally active iron chelators CP20, CP502, CP509, and ICL670)
- RN 243987-44-2 CAPLUS
- CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



- RN 793651-87-3 CAPLUS
- CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,N,1,6-tetramethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 19 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2004:723754 CAPLUS

DOCUMENT NUMBER: 141:391791

TITLE: Growth inhibition dependent on reactive oxygen species generated by C9-UK-2A, a derivative of the antifungal antibiotic UK-2A, in *Saccharomyces cerevisiae*


AUTHOR(S): Fujita, Kenichi; Tani, Kazunori; Usuki, Yoshinosuke; Tanaka, Toshio; Taniguchi, Makoto

10/580,011

CORPORATE SOURCE: Graduate School of Science, Osaka City University,
Osaka, 558-8585, Japan
SOURCE: Journal of Antibiotics (2004), 57(8), 511-517
CODEN: JANTAJ; ISSN: 0021-8820
PUBLISHER: Japan Antibiotics Research Association
DOCUMENT TYPE: Journal
LANGUAGE: English

AB UK-2A is a potent antifungal antibiotic and its structure is highly similar to that of antimycin A3 (AA). UK-2A and AA inhibit mitochondrial electron transport at complex III. C9-UK-2A, which has been prepared to improve the duration of the antifungal activity of UK-2A, shows durable fungicidal activities against various species of fungi and induces both membrane injury and the generation of cellular reactive oxygen species (ROS) against *Rhodotorula mucilaginosa* IFO 0001 cells. We found that C9-UK-2A inhibited the vegetative growth of *Saccharomyces cerevisiae* IFO 0203 cells accompanying cellular ROS generation in Sabouraud dextrose (SD) medium, which contained a fermentable carbon source. The ROS generation was completely restricted by pretreatment with a lipophilic antioxidant α -tocopherol. In addition, the pretreatment with the antioxidant protected against the growth inhibition induced by C9-UK-2A. C9-UK-2A also induced ROS generation in isolated mitochondria of the *S. cerevisiae* cells. The addition of both a complex I inhibitor rotenone and a complex II inhibitor thenoyltrifluoroacetone reduced ROS generation induced by C9-UK-2A in the whole cells and the isolated mitochondria. The addition of the inhibitors of complex III, AA or myxothiazol, or of complex IV, KCN, did not reduce ROS generation. These results suggest that C9-UK-2A induces ROS generation due to the blockade of electron flow at complex III, thereby inhibiting the growth of *S. cerevisiae* in SD medium.

IT 437651-11-1, C9-UK-2A
RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); BIOL (Biological study)
(growth inhibition dependent on reactive oxygen species generated by C9-UK-2A in *Saccharomyces cerevisiae*)
RN 437651-11-1 CAPLUS
CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-nonyl- (CA INDEX NAME)





REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 20 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2004:716288 CAPLUS
DOCUMENT NUMBER: 141:218924
TITLE: Antiviral agents containing nitrogen-containing heteroaromatic compounds
INVENTOR(S): Fuji, Masahiro; Matsushita, Shihaku; Mikamiyama, Hidenori
PATENT ASSIGNEE(S): Shionogi and Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 54 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

10/580,011

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 2004244320	A	20040902	JP 2003-32772	20030210
PRIORITY APPLN. INFO.:			JP 2003-32772	20030210
OTHER SOURCE(S):	MARPAT 141:218924			
GI				



AB The invention provides antiviral agents having HIV integrase inhibitory effects, characterized by containing I [G1 = (substituted) N,; G2 = (substituted) C; G3 = (substituted) N, C, O, S; R1 = (substituted) aryl, heteroaryl, cycloalkyl, cycloalkenyl, heterocycle; V1, V3 = (substituted) alkylene, alkenylene; V2 = (substituted) alkylene, alkenylene, etc.; X = O, S, NH; Y = hydroxy, mercapto, amino; Z = O, S, NH]. A compound II was prepared, and in vitro tested for its HIV integrase inhibitory effect. A capsule containing an active component 250 mg/capsule was also formulated.


IT 745803-24-1P 745803-26-3P

RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(antiviral agents having HIV integrase inhibitory effects containing nitrogen-containing heteroarom. compds.)


RN 745803-24-1 CAPLUS

CN 2-Pyridinecarboxamide, N-[(4-fluorophenyl)methyl]-3,4-dihydroxy- (CA INDEX NAME)



RN 745803-26-3 CAPLUS

CN 2-Pyridinecarboxamide, N-[2-(4-fluorophenyl)ethyl]-3,4-dihydroxy- (CA INDEX NAME)



L4 ANSWER 21 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:788612 CAPLUS

DOCUMENT NUMBER: 140:104367

TITLE: Improved high-performance liquid chromatographic method for the pharmacokinetic studies of a novel iron chelator, CP502, in rats

AUTHOR(S): Novakovic, Jasmina; Tesoro, Angelo; Spino, Michael; Thiessen, Jake

CORPORATE SOURCE: Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON, M5S 2S2, Can.

SOURCE: Journal of Chromatography, B: Analytical Technologies in the Biomedical and Life Sciences (2003), 796(1), 105-112

CODEN: JCBAAI; ISSN: 1570-0232

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An improved reverse-phase high-performance liquid chromatog. method (RP-HPLC) for the determination of a novel iron chelator CP502 (1,6-dimethyl-3-hydroxy-4-(1H)-pyridinone-2-carboxy-(N-methyl)-amide hydrochloride) in rat plasma, urine and feces was developed and validated. The separation was performed on a polymeric column using a mobile phase composed of 1 mM ethylenediaminetetra-acetic acid disodium salt (EDTA), acetonitrile, methanol and methylene chloride. Separation of CP502 from plasma, urine or feces endogenous compds. was achieved by gradient elution. Retention times of CP502 and its major metabolite (glucuronide) were about 13 and 4 min, resp. The method was validated in terms of limit of detection (LOD), limit of quantification (LOQ), selectivity (endogenous from plasma, urine or feces), linearity, extraction recovery, robustness (column selection, mobile phase composition, detection mode, internal standard

(IS) selection, analyte stability), day-to-day reproducibility and system suitability (repeatability, peak symmetry and resolution). The method is applicable to bioavailability and pharmacokinetic studies of CP502 in rats.


IT 243987-44-2, CP502

RL: PKT (Pharmacokinetics); BIOL (Biological study)

(improved high-performance liquid chromatog. method for pharmacokinetic studies of a novel iron chelator, CP502, in rats)

RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



10/580,011

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 22 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2003:280587 CAPLUS

DOCUMENT NUMBER: 139:242791

TITLE: UK-2A, B, C, and D, novel antifungal antibiotics from *Streptomyces* sp. 517-02. VIII. Reactive oxygen species generated by C9-UK-2A, a derivative of UK-2A, in *Rhodotorula mucilaginosa* IFO 0001

AUTHOR(S): Tani, Kazunori; Usuki, Yoshinosuke; Fujita, Ken-Ichi; Taniguchi, Makoto

CORPORATE SOURCE: Graduate School of Science, Osaka City University, Osaka, 558-8585, Japan

SOURCE: Journal of Antibiotics (2003), 56(3), 314-317
CODEN: JANTAJ; ISSN: 0021-8820

PUBLISHER: Japan Antibiotics Research Association

DOCUMENT TYPE: Journal

LANGUAGE: English


AB C9-UK-2A stimulated the generation of reactive oxygen species (ROS) in *R. mucilaginosa* in a dose- and time-dependent fashion and gradually decreased the number of CFU of *R. mucilaginosa*. Treatment with the lipophilic antioxidant α -tocopherol suppressed ROS generation caused by C9-UK-2A and almost completely stopped the decrease in cell viability, although viability did not recover to control levels. These results indicate that the antifungal activity of C9-UK-2A does not only depend on membrane injury and that ROS production alone does not fully explain the fungicidal effect of C9-UK-2A.

IT 437651-11-1, C9-UK-2A

RL: BSU (Biological study, unclassified); BIOL (Biological study)
(reactive oxygen species generated by the UK-2A derivative C9-UK-2A in *Rhodotorula mucilaginosa* IFO 0001)

RN 437651-11-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-nonyl- (CA INDEX NAME)



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 23 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:688565 CAPLUS

DOCUMENT NUMBER: 137:216878

TITLE: Preparation of 3-hydroxypyridin-4-ones as orally active iron(III) chelators.

INVENTOR(S): Hider, Robert Charles; Tilbrook, Gary Stuart; Liu, Zudong

PATENT ASSIGNEE(S): BTG International Limited, UK

SOURCE: U.S., 29 pp., Cont.-in-part of U.S. Ser. No. 437,211.
CODEN: USXXAM


DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6448273	B1	20020910	US 1999-451112	19991130
WO 9854138	A1	19981203	WO 1998-GB1517	19980526
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
US 6335353	B1	20020101	US 1999-437211	19991110
US 2002068758	A1	20020606	US 2001-944113	20010904
US 6506911	B2	20030114		
PRIORITY APPLN. INFO.:			GB 1997-11093	A 19970529
			WO 1998-GB1517	W 19980526
			US 1999-437211	A2 19991110
			US 1999-451112	A3 19991130
OTHER SOURCE(S):			MARPAT 137:216878	
GI				




AB Title compds. [I; R = H, group removable by metabolism in vivo to provide the free OH compound; R1 = aliphatic hydrocarbon group (un)substituted by OH or a carboxylic acid ester, sulfo acid ester, alkoxy, aryloxy, aralkoxy ether; R3 = H, alkyl; R4 = H, alkyl, alkyl, R2; R2 = CONHR5, CH2NHCO-R5, SO2NHR5, CH2NHSO2R5, CR6R6OR7, (viii) CONHCOR5; R5 = H, optionally hydroxy, alkoxy, or aralkoxy substituted 3alkyl, aryl, aralkyl; R6 = H, alkyl, aryl, aralkyl; R7 = H, alkyl, aryl, aralkyl; with provisos], were prepared Thus, 2-methoxymethyl-3-benzyloxy-6-methylpyran-4(1H)-one, MeNH2, and NaOH in H2O/EtOH were heated at 70° in a sealed tube for 12 h to give 82% 1,6-dimethyl-2-methoxymethyl-3-benzyloxypyridin-4(1H)-one hydrochloride. I at 150-450 µmol/kg orally in rats gave 6.3-73.5% Fe mobilization.

IT 216581-66-7P 216581-68-9P 216581-69-0P
216581-72-5P 216581-74-7P
RL: PAC (Pharmacological activity); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of 3-hydroxypyridin-4-ones as orally active iron(III) chelators)

RN 216581-66-7 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)

10/580,011




● HCl

RN 216581-68-9 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-N-(1-methylethyl)-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl


RN 216581-69-0 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-methoxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 216581-72-5 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-hydroxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)

10/580,011



RN 216581-74-7 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,N,1,6-tetramethyl-4-oxo-,
monohydrochloride (9CI) (CA INDEX NAME)



REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 24 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2002:570719 CAPLUS
DOCUMENT NUMBER: 137:125089
TITLE: Processes for manufacture of 3-hydroxy-N,1,6-trialkyl-
4-oxo-1,4-dihydropyridine-2-carboxamides
INVENTOR(S): Tam, Tim F.; Li, Wanren
PATENT ASSIGNEE(S): Apotex, Inc., Can.
SOURCE: U.S., 7 pp.
CODEN: USXXAM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6426418	B1	20020730	US 2001-985269	20011102
US 6472532	B1	20021029	US 2002-100133	20020319
US 6476229	B1	20021105	US 2002-100107	20020319
WO 2003037867	A1	20030508	WO 2002-CA1623	20021030

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,
UA, UG, US, UZ, VN, YU, ZA, ZM, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2002336835	A1	20030512	AU 2002-336835	20021030
EP 1440061	A1	20040728	EP 2002-771927	20021030
EP 1440061	B1	20070411		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK

BR 2002013814	A	20041019	BR 2002-13814	20021030
CN 1578769	A	20050209	CN 2002-821567	20021030
AT 359272	T	20070515	AT 2002-771927	20021030
ES 2284925	T3	20071116	ES 2002-2771927	20021030
IN 2004MN00249	A	20051118	IN 2004-MN249	20040427
MX 2004PA04063	A	20040708	MX 2004-PA4063	20040429
IN 2004MN00666	A	20060707	IN 2004-MN666	20041122

PRIORITY APPLN. INFO.: US 2001-985269 A3 20011102
WO 2002-CA1623 W 20021030

OTHER SOURCE(S): CASREACT 137:125089; MARPAT 137:125089
GI


* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A novel process for preparation of the title compds. I useful as orally active iron chelators comprises TEMPO oxidation of 3-O-protected-2-hydroxymethyl-6-alkyl-4H-pyran-4-one (III; R1 = H, lower alkyl; R4 = H, lower alkyl, lower alkoxy; R5 = H, alc. protective group) to 3-O-protected-6-alkyl-4-oxo-4H-pyran-2-carboxylic acid (II). Reaction of II with methylamine and 1,1'-carbonyldiimidazole in an inert solvent affords 3-O-protected-N,1,6-trialkyl-4-oxo-1,4-dihydropyridine-2-carboxamide, which is deprotected to give I.

IT 243987-44-2P 444103-12-2P, N,N-Diethyl-3-hydroxy-1,6-dimethyl-4-oxo-1,4-dihydropyridine-2-carboxamide
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of 3-hydroxy-N,1,6-trialkyl-4-oxo-1,4-dihydropyridine-2-carboxamides)


RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



RN 444103-12-2 CAPLUS

CN 2-Pyridinecarboxamide, N,N-diethyl-1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 25 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:312731 CAPLUS

DOCUMENT NUMBER: 137:272811

TITLE: Human immunodeficiency virus type I replication inhibition by the bidentate iron chelators CP502 and CP511 is caused by proliferation inhibition and the onset of apoptosis

AUTHOR(S): Georgiou, N. A.; van der Bruggen, T.; Oudshoorn, M.; Hider, R. C.; Marx, J. J. M.; van Asbeck, B. S.

CORPORATE SOURCE: Department of Internal Medicine and Eijkman-Winkler Institute for Microbiology Diseases and Inflammation, University Medical Center Utrecht, Utrecht, CX 3584, Neth.

SOURCE: European Journal of Clinical Investigation (2002), 32(Suppl. 1), 91-96

CODEN: EJCIB8; ISSN: 0014-2972

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Background: The iron chelators deferoxamine (DF) and deferiprone (CP20) have been shown to inhibit human immunodeficiency virus type 1 (HIV-1) replication in human peripheral blood lymphocytes (PBL). The orally active bidentate chelators CP502 and CP511, which also belong to the 3-hydroxypyridin-4-one family, but with higher affinities for iron than CP20, were monitored for their antiviral properties by checking for p24 antigen production and nuclear factor (NF)- κ B activation, and their ability to induce apoptosis. Materials and methods: Human PBLs were isolated from HIV-1 seroneg. donors and subsequently infected with HIV-1Ba-L for 2 h. After 5 days' incubation, HIV-1 replication was monitored by p24 antigen production. Cellular proliferation as well as caspase-3 activity were monitored in uninfected cells after a period of 5 days and after 1 day infection, resp. NF- κ B activity was also monitored by electromobility shift assays (EMSA) performed on nuclear exts. of Jurkat cells treated with the different chelators for 4 h. Results CP502 and CP511 decrease HIV-1 replication by decreasing cellular proliferation in a similar manner to DF and CP20. CP511 seemed to be more potent than either CP502 or CP20. Due to the reduction in cellular proliferation, there was an increase in caspase-3 activity after 24 h incubation. NF- κ B activity was not affected by any of the chelators. Conclusions: Iron chelators with high affinities for iron, which are under development for the treatment of iron overload, could contribute to the reduction of HIV-1 replication in infected patients by cellular proliferation inhibition rather than by a direct antiviral action.

IT 243987-44-2

RL: DMA (Drug mechanism of action); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)


(HIV type I replication inhibition by bidentate iron chelators CP502 and CP511 is caused by proliferation inhibition and onset of apoptosis

10/580,011

in human peripheral blood lymphocytes)

RN 243987-44-2 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 26 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2002:262139 CAPLUS

DOCUMENT NUMBER: 137:30441

TITLE: UK-2A, B, C, and D, novel antifungal antibiotics from *Streptomyces* sp. 517-02: VII. Membrane injury induced by C9-UK-2A, a derivative of UK-2A, in *Rhodotorula mucilaginosa* IFO 0001

AUTHOR(S): Tani, Kazunori; Usuki, Yoshinosuke; Motoba, Kazuhiko; Fujita, Ken-Ichi; Taniguchi, Makoto

CORPORATE SOURCE: Graduate School of Science, Osaka City University, Osaka, 558-8585, Japan


SOURCE: Journal of Antibiotics (2002), 55(3), 315-321
CODEN: JANTAJ; ISSN: 0021-8820

PUBLISHER: Japan Antibiotics Research Association

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



I

AB UK-2A is a potent antifungal antibiotic and its structure is highly similar to that of antimycin A3 (AA). UK-2A and AA inhibit mitochondrial electron transport at complex III. However, the antifungal activities of UK-2A and AA disappear after 48-h treatment. In an attempt to improve the duration of the antifungal activity of UK-2A, several UK-2A derivs. were prepared by substituting its nine-membered dilactone ring with an n-alkyl or an isoprenyl moiety. Among all the derivs. tested, C9-UK-2A (I) and C10-UK-2A showed the most potent and durable antifungal activities against a strict aerobic yeast, *Rhodotorula mucilaginosa* IFO 0001. I, in particular, continued to demonstrate its broad-spectrum antifungal activity after 120-h treatment. Therefore, we focused on I to further examine its mode of action against the yeast. Interestingly, I did not inhibit cellular respiration of the cells even at concns. greater than 100 µg/mL. I gradually induced the efflux of potassium ions from the cells. Moreover, I gradually induced the release of glucose from glucose-encapsulating liposomes. The patterns of efflux and release

10/580,011


induced by I were not as rapid as those seen with amphotericin B. These results suggest a membrane injury caused by I in *R. mucilaginosa* IFO 0001.

IT 267416-35-3, C8-UK-2A 321598-14-5 366791-61-9,
C16-UK-2A 366791-62-0, C12-UK-2A 366791-63-1, C4-UK-2A
366791-64-2 366791-65-3 437651-13-3, C10-UK-2A
437651-14-4, C11-UK-2A

RL: PAC (Pharmacological activity); BIOL (Biological study)
(activity of UK-2A and derivs. against *Rhodotorula mucilaginosa*)

RN 267416-35-3 CAPLUS


CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-octyl- (CA INDEX NAME)



RN 321598-14-5 CAPLUS


CN 2-Pyridinecarboxamide, N-[(2E)-3,7-dimethyl-2,6-octadienyl]-3-hydroxy-4-methoxy- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 366791-61-9 CAPLUS

CN 2-Pyridinecarboxamide, N-hexadecyl-3-hydroxy-4-methoxy- (CA INDEX NAME)



RN 366791-62-0 CAPLUS

CN 2-Pyridinecarboxamide, N-dodecyl-3-hydroxy-4-methoxy- (CA INDEX NAME)



RN 366791-63-1 CAPLUS


CN 2-Pyridinecarboxamide, N-butyl-3-hydroxy-4-methoxy- (CA INDEX NAME)

10/580,011

(membrane injury induced by the Streptomyces antifungal antibiotic
UK-2A derivative C9-UK-2A in Rhodotorula mucilaginosa IFO 0001)

RN 437651-11-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-nonyl- (CA INDEX NAME)



REFERENCE COUNT: 15 THERE ARE 15 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 27 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:557166 CAPLUS

DOCUMENT NUMBER: 135:300904

TITLE: UK-2A, B, C and D, novel antifungal antibiotics from
Streptomyces sp. 517-02. VI (1). Structure-activity
relationships of UK-2A

AUTHOR(S): Usuki, Yoshinosuke; Tani, Kazunori; Fujita, Ken-Ichi;
Taniguchi, Makoto

CORPORATE SOURCE: Graduate School of Science, Osaka City University,
Osaka, 558-8585, Japan

SOURCE: Journal of Antibiotics (2001), 54(7), 600-602
CODEN: JANTAJ; ISSN: 0021-8820

PUBLISHER: Japan Antibiotics Research Association

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The synthesis of UK-2A analogs, where the nine-membered dilactone residue
was replaced by several alkyl or isoprenyl moieties, and their biol.
effects were studied. All the tested compds., such as UK-2A, AA, and
their derivs., did not show any growth inhibitory activity against both
Gram-neg. and Gram-pos. bacteria up to 100µg/mL. Salicylic acid moiety
or pyridinecarboxylic acid moiety plus a hydrophobic structure is at least
necessary for expression of antifungal action. The 9-membered dilactone
ring moiety itself is not essential for the antimicrobial activity, and
C8-alkyl group is flexible and hydrophobic that makes C8-UK-2A interact
the binding domain to prevent yeasts and filamentous fungi from growing.
The decrease in activity of isoprenylated UK-2A derivs. was due to a loss
of flexibility, which interferes in their taking active conformations. AA
had strong cytotoxicity against porcine renal proximal tubule LLC-PK1
cells and other types of cultured cells compared to UK-2A. The inhibitory
of UK-2A and AA for the uncoupler stimulated respiration of bovine heart
submitochondrial particles was examined C8-3MeOSA showed comparably high
inhibitory activity similar to C8-AA and AA, although its antimicrobial
activities were weaker than those were. The mode of action of C8-UK-2A
would be different from that of UK-2A.

IT 267416-35-3 321598-14-5 366791-61-9
366791-62-0 366791-63-1 366791-64-2
366791-65-3


RL: BAC (Biological activity or effector, except adverse); BSU (Biological
study, unclassified); BIOL (Biological study)

(UK-2A, B, C and D, novel antifungal antibiotics from Streptomyces sp.
517-02. VI (1). Structure-activity relationships of UK-2A)

RN 267416-35-3 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-octyl- (CA INDEX NAME)


10/580,011



RN 321598-14-5 CAPLUS


CN 2-Pyridinecarboxamide, N-[(2E)-3,7-dimethyl-2,6-octadienyl]-3-hydroxy-4-methoxy- (9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 366791-61-9 CAPLUS

CN 2-Pyridinecarboxamide, N-hexadecyl-3-hydroxy-4-methoxy- (CA INDEX NAME)




RN 366791-62-0 CAPLUS

CN 2-Pyridinecarboxamide, N-dodecyl-3-hydroxy-4-methoxy- (CA INDEX NAME)



RN 366791-63-1 CAPLUS

CN 2-Pyridinecarboxamide, N-butyl-3-hydroxy-4-methoxy- (CA INDEX NAME)




RN 366791-64-2 CAPLUS


CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[(2E,6E)-3,7,11-trimethyl-2,6,10-dodecatrienyl]- (9CI) (CA INDEX NAME)

10/580,011

Double bond geometry as shown.



RN 366791-65-3 CAPLUS
CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(3-methyl-2-butenyl)- (9CI)
(CA INDEX NAME)



REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 28 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:507680 CAPLUS

DOCUMENT NUMBER: 135:92548

TITLE: Preparation of hydroxypicolinic acid derivatives for
agrochemical and pharmaceutical use as fungicides

INVENTOR(S): Bacque, Eric; Barriere, Jean-Claude; Vors,
Jean-Pierre; Nieto-Roman, Francisco; Villier, Alain

PATENT ASSIGNEE(S): Aventis CropScience SA, Fr.; Aventis Pharma S.A.

SOURCE: PCT Int. Appl., 100 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: French



FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001049667	A1	20010712	WO 2001-FR44	20010108
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2803592	A1	20010713	FR 2000-140	20000106
AT 340160	T	20061015	AT 2001-903877	20010105
ES 2272440	T3	20070501	ES 2001-1903877	20010105
CA 2396306	A1	20010712	CA 2001-2396306	20010108
EP 1248771	A1	20021016	EP 2001-903885	20010108
EP 1248771	B1	20060503		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			

10/580,011

BR 2001007425	A	20021203	BR 2001-7425	20010108
JP 2003519215	T	20030617	JP 2001-550207	20010108
HU 2003000139	A2	20030628	HU 2003-139	20010108
AT 325098	T	20060615	AT 2001-903885	20010108
IN 2002MN00517	A	20060505	IN 2002-MN517	20020422
ZA 2002003830	A	20031126	ZA 2002-3830	20020514
MX 2002PA06671	A	20021023	MX 2002-PA6671	20020704
US 2006040995	A1	20060223	US 2002-169855	20020708
PRIORITY APPLN. INFO.:			FR 2000-140	A 20000106
OTHER SOURCE(S):	MARPAT 135:92548		WO 2001-FR44	W 20010108
GI				



AB Hydroxypicolinic acid derivs., such as I [Q1 = O, imino, aminoimino; Q2 = alkyloxy, alkylthio, cycloalkyloxy, cycloalkylthio, amino, etc.; Y = H, OH, NH₂, N₃, CN, NO₂, alkyloxy, alkylthio, acylamino, etc.; Z = H, alkyl, aryl, allyl, propargyl, cycloalkyl, etc.; n = 0, 1], were prepared for agrochem. and pharmaceutical use as fungicides. Thus, picolinamide II was prepared by amidation of 3-hydroxy-4-methoxypyridine-2-carboxylic acid with 4-phenoxyaniline using 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride in pyridine at 75-85° for 1-2 h. Fungicidal biol. testing data for the prepared hydroxypicolinates was not presented.


IT 267415-60-1P 267415-69-0P 267415-77-0P
 267415-89-4P 267416-16-0P 267416-48-8P
 267416-59-1P 267416-63-7P 313643-54-8P
 313643-77-5P 313643-78-6P 348633-77-2P
 348633-78-3P 348633-79-4P 348633-80-7P
 348633-81-8P 348634-44-6P 348634-45-7P
 348634-47-9P 348634-48-0P 348634-49-1P
 348634-50-4P 348634-51-5P 348634-52-6P
 348634-69-5P 348634-70-8P 348634-71-9P
 348634-72-0P 348634-73-1P 348634-74-2P
 348634-75-3P 348634-76-4P 348634-77-5P
 348634-80-0P 348634-81-1P 348634-82-2P
 348634-83-3P 348634-84-4P 348634-85-5P
 348634-86-6P 348634-87-7P 348634-88-8P
 348634-89-9P 348634-90-2P 348634-91-3P
 348634-92-4P 348634-93-5P 348634-99-1P
 348635-00-7P 348635-01-8P 348635-02-9P
 348635-03-0P 348635-21-2P

RL: AGR (Agricultural use); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of hydroxypicolinic acid derivs. for agrochem. and pharmaceutical use as fungicides)

RN 267415-60-1 CAPLUS
 CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(4-phenoxyphenyl)- (CA INDEX NAME)


10/580,011

INDEX NAME)



RN 348635-21-2 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(4-methoxyphenyl)- (CA INDEX NAME)



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 29 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:507679 CAPLUS

DOCUMENT NUMBER: 135:92547

TITLE: Preparation of picolinic acid derivs. for agrochemical and therapeutic use as fungicides

INVENTOR(S): Nieto-Roman, Francisco; Vors, Jean-Pierre; Villier, Alain; Lachaise, Helene; Mousques, Adeline; Hartmann, Benoit; Hutin, Pierre; Molina, Jose Lorenzo; Muller, Benoit

PATENT ASSIGNEE(S): Aventis CropScience SA, Fr.

SOURCE: PCT Int. Appl., 121 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: French



FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001049666	A1	20010712	WO 2001-FR33	20010105
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
FR 2803592	A1	20010713	FR 2000-140	20000106
CA 2396299	A1	20010712	CA 2001-2396299	20010105
BR 2001007241	A	20020709	BR 2001-7241	20010105
EP 1244627	A1	20021002	EP 2001-903877	20010105
EP 1244627	B1	20060920		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			

10/580,011

HU 2002003958	A2	20030328	HU 2002-3958	20010105
JP 2003519214	T	20030617	JP 2001-550206	20010105
AT 340160	T	20061015	AT 2001-903877	20010105
ES 2272440	T3	20070501	ES 2001-1903877	20010105
AT 325098	T	20060615	AT 2001-903885	20010108
IN 2002MN00572	A	20040228	IN 2002-MN572	20020506
ZA 2002003830	A	20031126	ZA 2002-3830	20020514
BG 106834	A	20030131	BG 2002-106834	20020618
MX 2002PA06616	A	20021023	MX 2002-PA6616	20020702
US 2003191113	A1	20031009	US 2002-181842	20020708
PRIORITY APPLN. INFO.:			FR 2000-140	A 20000106
			WO 2001-FR33	W 20010105
OTHER SOURCE(S):	MARPAT 135:92547			
GI				




AB Picolinic acid derivs., such as I [Q1 = O, imino, aminoimino; Q2 = alkyloxy, alkylthio, cycloalkyloxy, cycloalkylthio, amino, etc.; Y = H, OH, NH₂, N₃, CN, NO₂, alkyloxy, alkylthio, acylamino, etc.; X1, X2 = H, OH, SH, NO₂, SCN, N₃, CN, halogen, alkyl, alkoxy, alkylthio, etc.; Z = H, alkyl, aryl, allyl, propargyl, cycloalkyl, etc.; n = 0, 1], were prepared for agrochem. use against plant fungal pathogens and pharmaceutical use as fungicides. Thus, picolinamide II was prepared by amidation of 3-hydroxy-4-methoxypyridine-2-carboxylic acid with 4-phenoxyaniline using 1-hydroxybenzotriazole and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride in pyridine at 85° for 2 h. The prepared picolinic acid derivs. were tested for activity against fungal strains, such as *Alternaria brassicae* and *Septoria nodorum*.

IT 348634-44-6P 348634-45-7P 348634-47-9P
 348634-48-0P 348634-49-1P 348634-50-4P
 348634-51-5P 348634-52-6P 348634-69-5P
 348634-70-8P 348634-71-9P 348634-72-0P
 348634-73-1P

RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of picolinic acid derivs. for agrochem. and therapeutic use as fungicides)


RN 348634-44-6 CAPLUS

CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-(4-phenoxyphenyl)- (CA INDEX NAME)




RN 348634-45-7 CAPLUS

CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-[4-(4-methylphenoxy)phenyl]- (CA



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 30 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 2001:322258 CAPLUS
 DOCUMENT NUMBER: 135:76762
 TITLE: Synthesis of 2-amido-3-hydroxypyridin-4(1H)-ones:
 novel iron chelators with enhanced pFe³⁺ values
 AUTHOR(S): Liu, Zu D.; Piyamongkol, S.; Liu, Ding Y.; Khodr,
 Hicham H.; Lu, Shu L.; Hider, Robert C.
 CORPORATE SOURCE: Department of Pharmacy, King's College London, London,
 SE1 8WA, UK
 SOURCE: Bioorganic & Medicinal Chemistry (2001), 9(3), 563-573
 CODEN: BMECEP; ISSN: 0968-0896
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 135:76762
 GI




AB The synthesis of a range of 2-amido-3-hydroxypyridin-4-ones as bidentate iron(III) chelators with potential for oral administration is described. The pK_a values of the ligands together with the stability consts. of their iron(III) complexes have been determined. Results indicate that the introduction of an amido substituent at the 2-position leads to an appreciable enhancement of the pFe³⁺ values. The ability of these novel 3-hydroxypyridin-4-ones to facilitate iron excretion in bile was investigated using a 59Fe-ferritin loaded rat model. The optimal effect was observed with the N-methylamido derivative I, which has an associated pFe³⁺ value of 21.7, more than two orders of magnitude higher than that of deferiprone (1,2-dimethyl-3-hydroxypyridin-4-one) (pFe³⁺ = 19.4). Dose-response studies suggest that chelators with high pFe³⁺ values scavenge iron more effectively at lower doses when compared with simple dialkyl-substituted hydroxypyridinones.

IT 216581-66-7P 216581-68-9P 216581-69-0P
 216581-72-5P 347393-49-1P 347393-50-4P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
 (preparation of 2-amido-3-hydroxypyridin-4(1H)-ones as iron chelators with enhanced pFe³⁺ values)

RN 216581-66-7 CAPLUS
 CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-,

10/580,011


monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-68-9 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-N-(1-methylethyl)-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-69-0 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-methoxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

RN 216581-72-5 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-hydroxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)


10/580,011



● HCl

RN 347393-49-1 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo-N-(phenylmethyl)-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 347393-50-4 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-4-oxo-N-phenyl-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

IT 216581-66-7D, complexes with iron 216581-68-9D,
complexes with iron 216581-69-0D, complexes with iron
216581-72-5D, complexes with iron


RL: PRP (Properties)

(preparation of 2-amido-3-hydroxypyridin-4(1H)-ones as iron chelators with enhanced pFe³⁺ values)

RN 216581-66-7 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)

10/580,011




● HCl

RN 216581-68-9 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-N-(1-methylethyl)-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl


RN 216581-69-0 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-methoxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-72-5 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-hydroxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)


10/580,011



● HCl


REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 31 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2001:279242 CAPLUS
DOCUMENT NUMBER: 135:92524
TITLE: Novel synthetic approach to 2-(1'-hydroxyalkyl)- and 2-amido-3-hydroxypyridin-4-ones
AUTHOR(S): Piyamongkol, S.; Liu, Z. D.; Hider, R. C.
CORPORATE SOURCE: Department of Pharmacy, King's College London, London, SE1 9NN, UK
SOURCE: Tetrahedron (2001), 57(16), 3479-3486
CODEN: TETRAB; ISSN: 0040-4020
PUBLISHER: Elsevier Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 135:92524
AB Novel methods for the synthesis of high pFe³⁺ iron chelators, 3,4-dihydroxy-2-(hydroxymethyl)pyridinium salts and 2-(aminocarbonyl)-3,4-dihydroxypyridinium compds., were reported. The products are obtained, via N-oxide intermediates, from either maltol or ethyl maltol. Iron-chelating properties were evaluated for 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-2-pyridinecarboxamide and 1,4-dihydro-3-hydroxy-N,6-dimethyl-4-oxo-2-pyridinecarboxamide.
IT 243987-44-2, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-2-pyridinecarboxamide 349141-34-0
RL: PRP (Properties)
(iron-chelating properties of 1,4-dihydro-3-hydroxy-4-oxo-pyridinecarboxamides)
RN 243987-44-2 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)




RN 349141-34-0 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,6-dimethyl-4-oxo- (CA INDEX NAME)

10/580,011




IT 349141-35-1P 349141-36-2P 349141-37-3P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of dihydroxy(hydroxymethyl)pyridinium compds. and
(aminocarbonyl)dihydroxypyridinium compds.)
RN 349141-35-1 CAPLUS
CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-methyl-, monohydrochloride (9CI)
(CA INDEX NAME)




● HCl

RN 349141-36-2 CAPLUS
CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-(2-hydroxyethyl)-,
monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 349141-37-3 CAPLUS
CN 2-Pyridinecarboxamide, 3,4-dihydroxy-N-phenyl-, monohydrochloride (9CI)
(CA INDEX NAME)



● HCl

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 32 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:152650 CAPLUS

DOCUMENT NUMBER: 134:207831

TITLE: Preparation, composition and use of heterocyclic aromatic amides as fungicides

INVENTOR(S): Ricks, Michael John; Dent, William Hunter, III; Rogers, Richard Brewer; Yao, Chenglin; Nader, Bassam Salim; Miesel, John Louis; Fitzpatrick, Gina Marie; Meyer, Kevin Gerald; Niyaz, Noormohamed Mohamed; Morrison, Irene Mae; Henry, Matthew James; Adamski, Butz Jenifer Lynn; Gajewski, Robert Peter

PATENT ASSIGNEE(S): Dow Agrosciences LLC, USA

SOURCE: PCT Int. Appl., 200 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001014339	A2	20010301	WO 2000-US21523	20000804
WO 2001014339	A3	20011115		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6521622	B1	20030218	US 2000-620662	20000720
CA 2376275	A1	20010301	CA 2000-2376275	20000804
AU 200065267	A	20010319	AU 2000-65267	20000804
AU 778108	B2	20041118		
US 6355660	B1	20020312	US 2000-632930	20000804
EP 1204643	A2	20020515	EP 2000-952599	20000804
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
EP 1234823	A2	20020828	EP 2002-9583	20000804
EP 1234823	A3	20030618		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
EP 1234824	A1	20020828	EP 2002-9584	20000804
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
EP 1234825	A2	20020828	EP 2002-9585	20000804

EP 1234825	A3	20030618		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
EP 1234826	A2	20020828	EP 2002-9586	20000804
EP 1234826	A3	20030618		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
EP 1234827	A2	20020828	EP 2002-9590	20000804
EP 1234827	A3	20030618		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
TR 200200409	T2	20030321	TR 2002-409	20000804
BR 2000013469	A	20030429	BR 2000-13469	20000804
HU 2003000924	A2	20030828	HU 2003-924	20000804
HU 2003000924	A3	20031028		
JP 2003527324	T	20030916	JP 2001-518428	20000804
EP 1486489	A2	20041215	EP 2004-22082	20000804
EP 1486489	A3	20050511		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1493733	A2	20050105	EP 2004-22081	20000804
EP 1493733	A3	20050126		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
US 2002177578	A1	20021128	US 2001-22413	20011213
US 2003018052	A1	20030123	US 2001-22207	20011213
US 2003018012	A1	20030123	US 2001-22511	20011213
US 6706740	B2	20040316		
US 2003022902	A1	20030130	US 2001-22483	20011213
US 2003022903	A1	20030130	US 2001-23497	20011213
ZA 2002000435	A	20030117	ZA 2002-435	20020117
US 2004034025	A1	20040219	US 2002-307844	20021202
US 7034035	B2	20060425		
US 2004048864	A1	20040311	US 2002-307710	20021202
US 6927225	B2	20050809		
US 39991	E1	20080101	US 2003-647172	20030822
PRIORITY APPLN. INFO.:			US 1999-149977P	P 19990820
			US 1999-150248P	P 19990823
			US 2000-620662	A 20000720
			US 1999-144676P	P 19990720
			EP 2000-952599	A3 20000804
			US 2000-632930	A3 20000804
			WO 2000-US21523	W 20000804
OTHER SOURCE(S):	MARPAT 134:207831			
GI				

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Title compds. [I; wherein X1-X4 independently = O, S, NR1, N, CR2, bond; R1 = H, C1-3 alkyl, C2-3 alkenyl, C2-3 alkynyl, OH, CHF2, C1-4 alkoxy; R2 = H, F, Cl, Br, CN, OH, C1-3 alkyl, C1-3 haloalkyl cyclopropyl, C1-3 alkoxy; Z = O, S, NOH, NOR3; R3 = C1-3 alkyl; A = C1-14 alkyl, C1-14 alkynyl, C1-14 cycloalkyl, aryl, heteroaryl, Q; M = H, Si(t-Bu)Me2, Si(Ph)Me2, SiEt3, CZR4, SO2R5; R4 = H, C1-6 alkyl, C2-6 alkenyl, C2-6 alkynyl; R5 = aryl, heteroaryl, C1-6 alkyl, C2-6 alkenyl, C3-6 alkenyl, C3-6 alkynyl, C3-6 cycloalkyl; X, Y independently = O, S; W = O, CH2, bond; R = C1-8 alkyl, C2-8 alkenyl, C2-8 alkynyl, C3-8 cycloalkyl, aryl, heteroaryl; R11 = H, C1-3 alkyl, C2-5 alkenyl, C2-5 alkynyl; R10 = H, R, OR, OCOR, OCOOR; R8, R9 independently = H, C1-6 alkyl, C2-6 alkenyl; R6, R7 independently = H, C1-6 alkyl, C2-6 alkenyl, C2-5 alkynyl, C3-6 cycloalkyl] are prepared as

10/580,011


fungicides involving application methods of effective usage of title compds. to control fungi, particularly plant pathogens and wood decaying fungi. The invention also encompasses hydrates, salts and complexes thereof. The title compound II was prepared and tested as fungicide.

IT 321598-36-1P 321599-09-1P

RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation and fungicidal activity of heterocyclic aromatic amides)

RN 321598-36-1 CAPLUS


CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(3,3,5,5-tetramethyl-4-oxocyclohexyl)- (CA INDEX NAME)



RN 321599-09-1 CAPLUS

CN L-Arabinonic acid, 2,5-dideoxy-2-(phenylmethyl)-, phenylmethyl ester, 3-(2-methylpropanoate), 4-ester with N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-O-(phenylmethyl)-L-serine (9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 166820-04-8P 267415-66-7P 267415-69-0P
267415-77-0P 267415-93-0P 313643-56-0P
313643-75-3P 313643-77-5P 313643-78-6P
321598-11-2P 321598-12-3P 321598-13-4P
321598-14-5P 321598-15-6P 321598-16-7P
321598-17-8P 321598-18-9P 321598-19-0P
321598-20-3P 321598-21-4P 321598-22-5P
321598-23-6P 321598-24-7P 321598-25-8P
321598-26-9P 321598-27-0P 321598-28-1P
321598-29-2P 321598-30-5P 321598-31-6P
321598-32-7P 321598-33-8P 321598-34-9P
321598-35-0P 321598-37-2P 321598-38-3P
321598-39-4P 321598-40-7P 321598-41-8P
321598-42-9P 321598-43-0P 321598-44-1P
321598-45-2P 321598-46-3P 321598-47-4P
321598-48-5P 321598-49-6P 321598-50-9P
321598-51-0P 321598-52-1P 321598-53-2P
321598-54-3P 321598-55-4P 321598-56-5P

10/580,011


Relative stereochemistry.



RN 321744-54-1 CAPLUS

CN 2-Pyridinecarboxamide, N-[(2R,3R)-4,5-dichloro-2,3,6,6-tetramethylcycloheptyl]-3-hydroxy-4-methoxy-, rel- (CA INDEX NAME)

Relative stereochemistry.




IT 321601-46-1

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation and fungicidal activity of heterocyclic aromatic amides)

RN 321601-46-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[(2R,3S)-2,3,6,6-tetramethyl-4-cyclohepten-1-yl]-, rel- (CA INDEX NAME)

Relative stereochemistry.



L4 ANSWER 33 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2001:113305 CAPLUS

DOCUMENT NUMBER: 134:320516

TITLE: Structure-activity investigation of the inhibition of 3-hydroxypyridin-4-ones on mammalian tyrosine hydroxylase

AUTHOR(S): Liu, Z. D.; Lockwood, M.; Rose, S.; Theobald, A. E.; Hider, R. C.

CORPORATE SOURCE: Department of Pharmacy, King's College London, London,


10/580,011

SOURCE: SE1 8WA, UK
Biochemical Pharmacology (2001), 61(3), 285-290
CODEN: BCPA6; ISSN: 0006-2952
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English


AB 3-Hydroxypyridin-4-ones are currently one of the main candidates for the development of orally active iron chelators. Small bidentate ligands tend to inhibit iron-containing metalloenzymes and therefore can cause undesirable side effects. A range of 3-hydroxypyridin-4-ones with different substituents at position 2 was selected for the investigation of the structure-activity relation between the chemical nature of the ligand and the inhibition of mammalian tyrosine hydroxylase. Results indicated that lipophilicity was the dominant factor in controlling the ability of this class of chelator to inhibit mammalian tyrosine hydroxylase. Ligands with hydrophilic substituents tended to be weak inhibitors. No significant correlation was found in this study between iron-binding affinity, extended substituent chain length, and enzyme inhibitory activity. In contrast, both the LogP values of the entire mol. and of the substituent segment correlated well with inhibitory activity.

IT 243987-44-2 336111-10-5
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(structure-activity relations of hydroxypyridinones as inhibitors of mammalian tyrosine hydroxylase)

RN 243987-44-2 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA INDEX NAME)



RN 336111-10-5 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-hydroxyethyl)-1,6-dimethyl-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 34 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2001:63978 CAPLUS
DOCUMENT NUMBER: 134:131431
TITLE: Fungicidal heterocyclic aromatic amides and their

10/580,011

INVENTOR(S): compositions, methods of use and preparation
Ricks, Michael John; Dent, William Hunter, III;
Rogers, Richard Brewer; Yao, Chenglin; Nader, Bassam
Salim; Miesel, John Louis; Fitzpatrick, Gina Marie;
Meyer, Kevin Gerald; Niyaz, Noormohamed Mohamed;
Morrison, Irene Mae; Gajewski, Robert Peter

PATENT ASSIGNEE(S): Dow Agrosiences LLC, USA

SOURCE: PCT Int. Appl., 159 pp.
CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001005769	A2	20010125	WO 2000-US19794	20000720
WO 2001005769	A3	20011122		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2374995	A1	20010125	CA 2000-2374995	20000720
EP 1196388	A2	20020417	EP 2000-950470	20000720
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO			
HU 2003001764	A2	20030929	HU 2003-1764	20000720
HU 2003001764	A3	20040628		
JP 2003528806	T	20030930	JP 2001-511430	20000720
BR 2000012615	A	20040330	BR 2000-12615	20000720
TR 200200587	T2	20041221	TR 2002-587	20000720
EP 1516874	A1	20050323	EP 2004-27006	20000720
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY			
EP 1516875	A1	20050323	EP 2004-27015	20000720
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY			
AU 780698	B2	20050414	AU 2000-63572	20000720
US 6355660	B1	20020312	US 2000-632930	20000804
US 2002177578	A1	20021128	US 2001-22413	20011213
US 2003018052	A1	20030123	US 2001-22207	20011213
US 2003018012	A1	20030123	US 2001-22511	20011213
US 6706740	B2	20040316		
US 2003022902	A1	20030130	US 2001-22483	20011213
US 2003022903	A1	20030130	US 2001-23497	20011213
KR 743262	B1	20070727	KR 2002-700677	20020116
ZA 2002000436	A	20040302	ZA 2002-436	20020117
US 2004034025	A1	20040219	US 2002-307844	20021202
US 7034035	B2	20060425		
US 2004048864	A1	20040311	US 2002-307710	20021202
US 6927225	B2	20050809		


PRIORITY APPLN. INFO.:

US 1999-144676P	P	19990720
US 1999-149977P	P	19990820
US 1999-150248P	P	19990823
EP 2000-950470	A3	20000720
US 2000-620662	A3	20000720
WO 2000-US19794	W	20000720
US 2000-632930	A3	20000804

OTHER SOURCE(S):

MARPAT 134:131431

GI




AB Title compds. I [W, X, Y, Z are selected from S, O, NR1, N, CR2 or bond and comprise a 5-6 membered (un)substituted heterocyclic ring; R1 = H, alkyl, alkenyl, alkynyl, OH, acyloxy, alkoxy, CHF2, cyclopropyl, or alkoxy; R2 = H, halo, CN, OH, alkyl, haloalkyl, cyclopropyl, alkoxy, haloalkoxy, etc.; G = O, S or NOR3 where R3 = H or alkyl; A = (un)substituted alkyl, alkenyl, alkynyl, cycloalkyl, unsatd. cycloalkyl, heterocycle, bi or tricyclic ring system which may contain heteroatoms, aryl or heteroaryl, etc.] bearing a hydroxy group adjacent to the amide functionality are prepared and disclosed as antifungal agents, particularly for plants. Thus, pyridinyl carboxamide II was prepared via amidation of 3-benzyloxy-6-bromo-4-methoxypyridin-2-carbonyl chloride with 4-(4-(trifluoromethylphenoxy)aniline with subsequent deprotection. The preferred fungicidal composition consists of a compound of formula I with a phytol acceptable carrier. Activity has been demonstrated against a variety of fungi, e.g., *Plasmopara viticola* (Downy Mildew of Grape), *Phytophthora infestans* (Late Blight of Tomato), and *Venturia inaequalis* (Apple Scab). I is both useful for eradication and prevention of fungal attack.

IT 321598-36-1P 321599-09-1P

RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation and fungicidal activity of heterocyclic aromatic amides)

RN 321598-36-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(3,3,5,5-tetramethyl-4-oxocyclohexyl)- (CA INDEX NAME)




RN 321599-09-1 CAPLUS

CN L-Arabinonic acid, 2,5-dideoxy-2-(phenylmethyl)-, phenylmethyl ester, 3-(2-methylpropanoate), 4-ester with N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-O-(phenylmethyl)-L-serine (9CI) (CA INDEX NAME)

Absolute stereochemistry.

10/580,011


Relative stereochemistry.



RN 321744-54-1 CAPLUS

CN 2-Pyridinecarboxamide, N-[(2R,3R)-4,5-dichloro-2,3,6,6-tetramethylcycloheptyl]-3-hydroxy-4-methoxy-, rel- (CA INDEX NAME)

Relative stereochemistry.




IT 321601-46-1

RL: RCT (Reactant); RACT (Reactant or reagent)
(preparation and fungicidal activity of heterocyclic aromatic amides)

RN 321601-46-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[(2R,3S)-2,3,6,6-tetramethyl-4-cyclohepten-1-yl]-, rel- (CA INDEX NAME)

Relative stereochemistry.



L4 ANSWER 35 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 2000:900620 CAPLUS

DOCUMENT NUMBER: 134:56577



TITLE: Pyridinecarboxamides and their use as plant protection agents

INVENTOR(S): Backhaus, Dirk; Jordan, Stephan; Boie, Christiane;
Schneider, Udo; Gayer, Herbert; Vaupel, Martin;
Mauler-Machnik, Astrid; Wachendorff-Neumann, Ulrike;
Kuck, Karl-Heinz

10/580,011

PATENT ASSIGNEE(S): Bayer A.-G., Germany
SOURCE: PCT Int. Appl., 63 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: German
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000076979	A1	20001221	WO 2000-EP4870	20000529
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
DE 19958166	A1	20001214	DE 1999-19958166	19991202
IN 2000MU00490	A	20050304	IN 2000-MU490	20000529
PRIORITY APPLN. INFO.:			DE 1999-19926174	A 19990609
			DE 1999-19958166	A 19991202
OTHER SOURCE(S):			MARPAT 134:56577	
GI				




AB Pyridinecarboxamides I [A = bond, (un)substituted alkylene, heteroalkylene; R1 = (un)substituted cycloalkyl, cycloalkenyl, aryl, heterocyclyl; R2 = H, acyl, alkoxycarbonyl] were prepared for use as agricultural fungicides. Thus, the amide II was obtained by amidation. II was ≥91% effective against Botrytis on beans at 500 g/ha.

IT 313643-52-6P
RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); RCT (Reactant); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); RACT (Reactant or reagent); USES (Uses)
(preparation of pyridinecarboxamides as agricultural fungicides)

RN 313643-52-6 CAPLUS

CN 2-Pyridinecarboxamide, N-[4-(2,4-dichlorophenoxy)phenyl]-3-hydroxy-4-methoxy- (CA INDEX NAME)




IT 267415-65-6P 267415-77-0P 267415-79-2P
 267415-86-1P 267416-59-1P 313643-54-8P
 313643-55-9P 313643-56-0P 313643-57-1P
 313643-58-2P 313643-59-3P 313643-60-6P
 313643-61-7P 313643-62-8P 313643-63-9P
 313643-64-0P 313643-65-1P 313643-66-2P
 313643-69-5P 313643-73-1P 313643-75-3P
 313643-76-4P 313643-77-5P 313643-78-6P
 313643-79-7P 313643-80-0P 313643-81-1P
 313643-82-2P 313643-83-3P 313643-84-4P
 313643-85-5P 313643-86-6P 313643-87-7P
 313643-88-8P 313643-89-9P 313643-90-2P
 313643-91-3P 313643-94-6P

RL: AGR (Agricultural use); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of pyridinecarboxamides as agricultural fungicides)


RN 267415-65-6 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[4-(phenylthio)phenyl]- (CA INDEX NAME)



RN 267415-77-0 CAPLUS


CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[4-[3-(trifluoromethyl)phenoxy]phenyl]- (CA INDEX NAME)



RN 267415-79-2 CAPLUS

CN 2-Pyridinecarboxamide, N-[3-chloro-4-(4-chlorophenoxy)phenyl]-3-hydroxy-4-methoxy- (CA INDEX NAME)


10/580,011



RN 313643-90-2 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-O-(phenylmethyl)-, methyl ester (CA INDEX NAME)


Absolute stereochemistry.



RN 313643-91-3 CAPLUS


CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, phenylmethyl ester (CA INDEX NAME)

Absolute stereochemistry.



RN 313643-94-6 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-[3-phenoxy-5-(trifluoromethyl)phenyl]- (CA INDEX NAME)



REFERENCE COUNT:

5

THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 36 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN


ACCESSION NUMBER: 2000:314676 CAPLUS

DOCUMENT NUMBER: 132:334362

TITLE: Preparation of picolinamide derivatives and pest controllers containing the same as the active

ingredient
 INVENTOR(S): Imamura, Keiichi; Mitomo, Kouichi; Yamada, Natsuko;
 Yamamoto, Kazumi; Teraoka, Takeshi; Sakanaka, Osamu;
 Kurihara, Hiroshi; Taniguchi, Makoto
 PATENT ASSIGNEE(S): Meiji Seika Kaisha, Ltd., Japan
 SOURCE: PCT Int. Appl., 98 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: Japanese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000026191	A1	20000511	WO 1999-JP6142	19991104
W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2353627	A1	20000511	CA 1999-2353627	19991104
EP 1134214	A1	20010919	EP 1999-954375	19991104
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
AU 771975	B2	20040408	AU 2000-10768	19991104
US 7183278	B1	20070227	US 2001-830923	20010809
PRIORITY APPLN. INFO.:			JP 1998-313688	A 19981104
			WO 1999-JP6142	W 19991104
OTHER SOURCE(S):		MARPAT 132:334362		
GI				



AB Described are novel compds. of general formula [I; wherein A is a bond or optionally substituted alkylene; R1 is one or more groups which may be the same or different from each other and are selected from among hydrogen, alkoxy and haloalkoxy; R2 is hydrogen, (substituted) benzyl, (substituted) alkyl or (substituted) alkanoyl; and R3 is hydrogen, (substituted) cycloalkyl, (substituted) cycloalkenyl, (substituted) aryl, or a (substituted) heterocyclic group, with the proviso that the cases wherein R1 is hydrogen, A is a free valency or methylene, and R3 is Ph or cyclohexyl or those wherein A is alkylene and R3 is hydrogen are excepted.], pest controllers such as plant fungicides, insecticides, and herbicides containing the same; and a process for the preparation of the compds.

Thus, a solution of 1.85 g 4-phenoxyaniline in 25 mL DMF was added dropwise to a suspension of 1.39 g 3-hydroxypicolinic acid, 1.95 g carbonyl diimidazole, and 30 mL DMF and stirred overnight to give 41% 3-hydroxy-4'-phenoxy-picolinanilide (II). II at 100 ppm protected 80-100% rice seedlings against Pyricularia oryzae.


IT 267415-60-1P 267415-61-2P 267415-62-3P
 267415-63-4P 267415-64-5P 267415-65-6P
 267415-66-7P 267415-67-8P 267415-68-9P

267415-69-0P 267415-70-3P 267415-71-4P
 267415-72-5P 267415-73-6P 267415-74-7P
 267415-75-8P 267415-76-9P 267415-77-0P
 267415-78-1P 267415-79-2P 267415-80-5P
 267415-81-6P 267415-82-7P 267415-83-8P
 267415-84-9P 267415-85-0P 267415-86-1P
 267415-87-2P 267415-88-3P 267415-89-4P
 267415-90-7P 267415-91-8P 267415-92-9P
 267415-93-0P 267415-94-1P 267415-95-2P
 267415-96-3P 267415-97-4P 267415-98-5P
 267415-99-6P 267416-02-4P 267416-03-5P
 267416-04-6P 267416-05-7P 267416-06-8P
 267416-07-9P 267416-08-0P 267416-09-1P
 267416-10-4P 267416-11-5P 267416-12-6P
 267416-13-7P 267416-15-9P 267416-16-0P
 267416-17-1P 267416-18-2P 267416-19-3P
 267416-21-7P 267416-22-8P 267416-24-0P
 267416-25-1P 267416-26-2P 267416-27-3P
 267416-28-4P 267416-29-5P 267416-30-8P
 267416-31-9P 267416-32-0P 267416-33-1P
 267416-34-2P 267416-35-3P 267416-36-4P
 267416-37-5P 267416-41-1P 267416-42-2P
 267416-48-8P 267416-49-9P 267416-50-2P
 267416-51-3P 267416-52-4P 267416-53-5P
 267416-54-6P 267416-55-7P 267416-56-8P
 267416-57-9P 267416-58-0P 267416-59-1P
 267416-60-4P 267416-61-5P 267416-62-6P
 267416-63-7P 267416-64-8P 267416-65-9P
 267416-66-0P 267416-67-1P 267416-68-2P
 267416-70-6P 267416-71-7P 267416-72-8P

RL: AGR (Agricultural use); BAC (Biological activity or effector, except
 adverse); BSU (Biological study, unclassified); SPN (Synthetic
 preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of picolinamide derivs. as pest controllers)


RN 267415-60-1 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(4-phenoxyphenyl)- (CA INDEX
 NAME)



RN 267415-61-2 CAPLUS

CN 2-Pyridinecarboxamide, N-[4-[4-(1,1-dimethylethyl)phenoxy]phenyl]-3-
 hydroxy-4-methoxy- (CA INDEX NAME)




RN 267415-62-3 CAPLUS

CN 2-Pyridinecarboxamide, 3-hydroxy-4-methoxy-N-(3-phenoxyphenyl)- (CA INDEX
 NAME)


10/580,011

CN 2-Pyridinecarboxamide, N-[4-(2,2-dimethylpropyl)phenyl]-3-hydroxy-4-methoxy- (CA INDEX NAME)




RN 267416-70-6 CAPLUS

CN 2-Pyridinecarboxamide, N-(3,4-dichlorophenyl)-3-hydroxy-4-methoxy- (CA INDEX NAME)




RN 267416-71-7 CAPLUS

CN 2-Pyridinecarboxamide, N-[4-(1,1-dimethylethyl)-2,6-dimethylphenyl]-3-hydroxy-4-methoxy- (CA INDEX NAME)



RN 267416-72-8 CAPLUS

CN 2-Pyridinecarboxamide, N-[2-chloro-4-(1,1-dimethylethyl)phenyl]-3-hydroxy-4-methoxy- (CA INDEX NAME)



REFERENCE COUNT: 19 THERE ARE 19 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 37 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1999:605552 CAPLUS

DOCUMENT NUMBER: 132:35975

TITLE: Synthesis of homorhamnojirimycins and related trihydroxypipelic acid derivatives via divergent bicyclic amino lactone intermediates: Inhibition of naringinase (L-rhamnosidase) and dTDP-rhamnose biosynthesis

10/580,011

AUTHOR(S): Shilvock, John P.; Wheatley, Joseph R.; Nash, Robert J.; Watson, Alison A.; Griffiths, Rhodri C.; Butters, Terry D.; Muller, Mathias; Watkin, David J.; Winkler, David A.; Fleet, George W. J.
CORPORATE SOURCE: Dyson Perrins Laboratory, Oxford University, Oxford, OX1 3QY, UK
SOURCE: Journal of the Chemical Society, Perkin Transactions 1: Organic and Bio-Organic Chemistry (1999), (19), 2735-2745
CODEN: JCPRB4; ISSN: 0300-922X
PUBLISHER: Royal Society of Chemistry
DOCUMENT TYPE: Journal
LANGUAGE: English

AB A series of homorhamnojirimycins and related compds. are prepared from two epimeric [2.2.2] bicyclic amino lactones via the 2-azidoheptono-1,5-lactone, itself derived from L-rhamnose. Aminolysis and deprotection of the bicyclic lactones provides an efficient route to trihydroxypipicolinic acid amide analogs of 5-epi-L-rhamnopyranose and L-rhamnopyranose. Some of the L-rhamnopyranose analogs display inhibitory activity against naringinase (L-rhamnosidase) and dTDP-rhamnose biosynthesis and are potentially useful as tools for investigating cell wall biosynthesis of Mycobacterium tuberculosis, the causative agent of tuberculosis. The synthesis of other homoiminosugar analogs including epi-homorhamnojirimycin (HRJ) is also reported. Methanolysis of the bicyclic lactone possessing a configuration corresponding to α -L-rhamnopyranose under basic conditions affords both α - and β -Me 2,6-iminoheptonates. Reduction and subsequent deprotection affords the 2,6-iminoheptitols, α -homorhamnojirimycin (α -HRJ) and β -homorhamnojirimycin (β -HRJ), potent inhibitors of L-rhamnosidase and α -galactosidase, resp. The crystal-structure determination of the bicyclic lactone is also reported.

IT 252358-34-2P 252358-38-6P 252358-39-7P
252358-40-0P 252358-44-4P 252358-45-5P
252358-46-6P


RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(synthesis of homorhamnojirimycins and related trihydroxypipicolinic acid derivs. and their inhibition of naringinase and dTDP-rhamnose biosynthesis)

RN 252358-34-2 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-N,6-dimethyl-, (2S,3S,4R,5S,6S) -
(CA INDEX NAME)

Absolute stereochemistry. Rotation (-).




RN 252358-38-6 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-N,6-dimethyl-, (2R,3S,4R,5S,6R) -
(CA INDEX NAME)

Absolute stereochemistry. Rotation (-).


10/580,011



RN 252358-39-7 CAPLUS

CN 2-Piperidinecarboxamide, N-butyl-3,4,5-trihydroxy-6-methyl-,
(2R,3S,4R,5S,6R)- (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 252358-40-0 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-methyl-N-(phenylmethyl)-,
(2R,3S,4R,5S,6R)- (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 252358-44-4 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-N,6-dimethyl-, (2R,3S,4R,5S,6S)-
(CA INDEX NAME)

Absolute stereochemistry. Rotation (+).




RN 252358-45-5 CAPLUS

10/580,011

CN 2-Piperidinecarboxamide, N-butyl-3,4,5-trihydroxy-6-methyl-,
(2R,3S,4R,5S,6S) - (CA INDEX NAME)


Absolute stereochemistry. Rotation (+).



RN 252358-46-6 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-methyl-N-(phenylmethyl)-,
(2R,3S,4R,5S,6S) - (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 38 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1999:486386 CAPLUS

DOCUMENT NUMBER: 131:222962

TITLE: Gradient ion-pair high-performance liquid
chromatographic method for analysis of
3-hydroxypyridin-4-one iron chelators

AUTHOR(S): Liu, Ding Y.; Liu, Zu D.; Lu, Shu L.; Hider, Robert C.

CORPORATE SOURCE: Department of Pharmacy, King's College London,
University of London, London, SW3 6LX, UK

SOURCE: Journal of Chromatography, B: Biomedical Sciences and
Applications (1999), 730(1), 135-139
CODEN: JCBEP; ISSN: 0378-4347

PUBLISHER: Elsevier Science B.V.


DOCUMENT TYPE: Journal

LANGUAGE: English


AB A gradient ion-pair HPLC separation of highly hydrophilic 3-hydroxypyridin-4-one (HPO) iron chelators is described. The separation of HPOs was performed using a reversed-phase polymer HPLC column (PLRP-S 100 A, 15+0.46 cm ID, 5 µm). The ion-pair buffer contained 1-heptanesulfonic acid (sodium salt) (5 mM) and the pH was adjusted to 2.0 using HCl. The gradient was 2%-35% CH₃CN in 20 min and post-run was followed for 5 min using 2% CH₃CN and 98% buffer. The flow-rate was 1 mL/min and the analytes were monitored at 280 nm. The retention times of 30 hydrophilic HPOs fell in the range of 10-18 min with sharp peak shapes, although these iron chelators possess various functional groups and distribution coeffs. The application of this HPLC method in the anal. of HPO chelators and their metabolites in rat bile and urine is described.

10/580,011

IT 243987-44-2 243987-45-3
RL: ANT (Analyte); ANST (Analytical study)
(ion pair HPLC for anal. of 3-hydroxypyridin-4-one iron chelators in
biol. fluids)
RN 243987-44-2 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo- (CA
INDEX NAME)






RN 243987-45-3 CAPLUS
CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-N-(1-
methylethyl)-4-oxo- (CA INDEX NAME)



REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 39 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 1999:19692 CAPLUS
DOCUMENT NUMBER: 130:168617
TITLE: UK-2A, B, C and D, novel antifungal antibiotics from
Streptomyces sp. 517-02 III. Absolute configuration of
an antifungal antibiotic, UK-2A, and consideration of
its conformation
AUTHOR(S): Shibata, Kozo; Hanafi, Muhammad; Fujii, Jyunko;
Sakanaka, Osamu; Iinuma, Katsuharu; Ueki, Masashi;
Taniguchi, Makoto
CORPORATE SOURCE: Faculty of Science, Osaka City University, Osaka,
558-8585, Japan
SOURCE: Journal of Antibiotics (1998), 51(12), 1113-1116
CODEN: JANTAJ; ISSN: 0021-8820
PUBLISHER: Japan Antibiotics Research Association
DOCUMENT TYPE: Journal
LANGUAGE: English
GI



AB The absolute configuration of UK-2A (I) was determined by the elucidation of the absolute configurations of butanolide II and the serine derivative III, the products of alkaline hydrolysis of I. The absolute configuration of UK-2A was found to be (+)-(2R,3R,4S,7S).

IT 166820-04-8P 166820-06-0P


RL: SPN (Synthetic preparation); PREP (Preparation)

(determination of the absolute configuration of UK-2A, an antifungal antibiotic)

RN 166820-04-8 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, methyl ester (CA INDEX NAME)


Absolute stereochemistry.



RN 166820-06-0 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:

6


THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

10/580,011

ACCESSION NUMBER: 1998:793129 CAPLUS
DOCUMENT NUMBER: 130:38296
TITLE: Preparation of 3-hydroxypyridin-4-ones as novel orally active iron(III) chelators, and their pharmaceutical formulations
INVENTOR(S): Hider, Robert Charles; Tilbrook, Gary Stuart; Liu, Zudong
PATENT ASSIGNEE(S): BTG International Limited, UK
SOURCE: PCT Int. Appl., 62 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9854138	A1	19981203	WO 1998-GB1517	19980526
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
CA 2287907	A1	19981203	CA 1998-2287907	19980526
AU 9875427	A	19981230	AU 1998-75427	19980526
AU 751600	B2	20020822		
EP 984934	A1	20000315	EP 1998-922968	19980526
EP 984934	B1	20030108		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002500663	T	20020108	JP 1999-500362	19980526
AT 230728	T	20030115	AT 1998-922968	19980526
ES 2187025	T3	20030516	ES 1998-922968	19980526
ZA 9804635	A	19991129	ZA 1998-4635	19980529
US 6335353	B1	20020101	US 1999-437211	19991110
MX 9910947	A	20000430	MX 1999-10947	19991126
US 6448273	B1	20020910	US 1999-451112	19991130
US 2002068758	A1	20020606	US 2001-944113	20010904
US 6506911	B2	20030114		
PRIORITY APPLN. INFO.:			GB 1997-11093	A 19970529
			WO 1998-GB1517	W 19980526
			US 1999-437211	A2 19991110
			US 1999-451112	A3 19991130

OTHER SOURCE(S): MARPAT 130:38296
GI



AB Novel 3-hydroxypyridin-4-ones I are provided, wherein R = H or a group that is removed by metabolism in vivo to provide the free hydroxy compound, R1 =

aliphatic hydrocarbon group (un)substituted by a hydroxy group or a carboxylic acid ester, sulfo acid ester or a C1-6alkoxy, C6-aryloxy or C7-10aralkoxy ether, R3 = H or C1-6alkyl; R4 = H, C1-6alkyl, C1-6alkyl, a group as described for R2 characterized in that R2 is selected from groups (i) -CONH-R5, (ii) -CH2NHCO-R5, (iii) -SO2NH-R5, (iv) -CH2NHSO2-R5, (v) -CR6R6OR7, (viii) -CONHCOR5, wherein R5 is selected from H and optionally hydroxy, alkoxy, or aralkoxy substituted C1-13alkyl, aryl and C7-13aralkyl, R6 is independently selected from H, C1-13alkyl, aryl and C7-13aralkyl, and R7 is selected from H, C1-13alkyl, aryl and C7-13aralkyl or a pharmaceutically acceptable salt of any such compound with the proviso that when R7 is H, R6 is not selected from aryl and with the proviso that the compound is not 1-ethyl-2-(1'-hydroxyethyl)-3-hydroxypyridin-4-one. Compds. I are orally active iron(III) chelators and are useful in the manufacture of a medicament for treatment of an iron-associated disease. Iron(III) mobilization efficacy assays of compds. I in rat are given. Pharmaceutical compns. containing I are claimed (3 examples). Processes for the preparation of I are also provided. Prepared intermediates include 8-oxo-4,8-dihydro-2-phenyl-4H-pyridino[3,2-d]-m-dioxins, 2-(1-hydroxyalkyl)-3-hydroxypyran-4(1H)-ones, and 3-benzyloxy-2-(2-thionothiazolidine-3-carbonyl)pyran-4(1H)-ones.

IT 216581-66-7P 216581-68-9P 216581-69-0P


216581-72-5P 216581-74-7P

RL: SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)

(preparation as prodrug for orally active iron(III) chelators)

RN 216581-66-7 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,1,6-trimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-68-9 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-1,6-dimethyl-N-(1-methylethyl)-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)




● HCl

10/580,011

RN 216581-69-0 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-methoxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-72-5 CAPLUS


CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-(2-hydroxyethyl)-1,6-dimethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

RN 216581-74-7 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N,N,1,6-tetramethyl-4-oxo-, monohydrochloride (9CI) (CA INDEX NAME)



● HCl

REFERENCE COUNT:

9

THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 41 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1998:773075 CAPLUS

DOCUMENT NUMBER: 130:110500

TITLE: Intermediates for incorporation of tetrahydroxypipelicolic acid analogs of α - and β -D-mannopyranose into combinatorial libraries: unexpected nanomolar-range hexosaminidase inhibitors. Synthesis of α - and β -homomannojirimycin

AUTHOR(S): Shilvock, John P.; Nash, Robert J.; Lloyd, Janet D.; Winters, Ana L.; Asano, Naoki; Fleet, George W. J.

CORPORATE SOURCE: Dyson Perrins Laboratory, Oxford University, Oxford, OX1 3QY, UK

SOURCE: Tetrahedron: Asymmetry (1998), 9(19), 3505-3516
CODEN: TASYE3; ISSN: 0957-4166

PUBLISHER: Elsevier Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

OTHER SOURCE(S): CASREACT 130:110500

AB Homoazasugars have the distinction as a class of natural products in that most of them have been synthesized before they were isolated. Syntheses of α - and β -homomannojirimycin rely on the stereoselective and chemo-selective sodium cyanoborohydride reduction of a [2.2.2] bicyclic imino lactone to give a single [2.2.2] bicyclic amino-lactone. Methanolysis under basic conditions is accompanied by efficient epimerization of the first formed α -amino-ester to the more stable β -amino-ester in which the 2,6-substituents are equatorial. Both the [2.2.2] bicyclic amino-lactone and the β -amino-ester are suitable intermediates for the incorporation of tetrahydroxypipelicolic acid derivs. into combinatorial libraries containing α - and β -C-glycosyl analogs of aza-D-mannopyranose, resp. Methylamides are shown to be specific and potent inhibitors of two β -N-acetylglucosaminidases but have no effect on an α -N-acetylgalactosaminidase. The synthesis of α - and β -manno-pipelicolic acids is also reported.


IT 219589-69-2P 219589-71-6P
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)

(intermediates for incorporation of tetrahydroxypipelicolic acid analogs of mannopyranose into combinatorial libraries)

RN 219589-69-2 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-, (2S,3R,4S,5R,6R) - (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).



RN 219589-71-6 CAPLUS


CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-N-methyl-, (2R,3R,4S,5R,6R) - (CA INDEX NAME)

Absolute stereochemistry. Rotation (+).



REFERENCE COUNT: 29 THERE ARE 29 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 42 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
 ACCESSION NUMBER: 1998:651994 CAPLUS
 DOCUMENT NUMBER: 130:3703
 TITLE: Total synthesis of the antifungal dilactones UK-2A and UK-3A: the determination of their relative and absolute configurations, analog synthesis and antifungal activities
 AUTHOR(S): Shimano, Masanao; Kamei, Noriyuki; Shibata, Tetsuo; Inoguchi, Kiyoshi; Itoh, Nobuko; Ikari, Takashi; Senda, Hisato
 CORPORATE SOURCE: Dep. Med. Chem. Mol. Design, Drug Discovery Res. Lab., Kaken Pharmaceutical Co., Ltd., Minami Kawara-cho, Yamashina-ku, Kyoto, 607-8042, Japan
 SOURCE: Tetrahedron (1998), 54(42), 12745-12774
 CODEN: TETRAB; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 130:3703
 GI



I

AB The synthesis of the antifungal dilactones (I), UK-2A (R = OMe) and UK-3A (R = H), is described. In addition to providing a workable synthetic route to these potent antifungal antibiotics, this has allowed us to determine the assignment of the relative and absolute configurations in the nine-membered ring. Furthermore, UK-2A analogs were also synthesized and evaluated for their antifungal activities and cytotoxic activities along with UK-2A, (2R, 3R, 4S, 7R)-UK-2A, UK-3A, (2R, 3R, 4S, 7R)-UK-3A, and antimycin A. The structural requirements for the selective cytotoxicity against yeasts and filamentous fungi will also be suggested.


IT 166820-04-8P 215874-80-9P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (synthesis, antifungal activity, cytotoxicity and absolute configuration of dilactones UK-2A and UK-3A)

10/580,011

RN 166820-04-8 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, methyl ester
(CA INDEX NAME)


Absolute stereochemistry.



RN 215874-80-9 CAPLUS

CN D-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, methyl ester
(CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 43 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1997:473732 CAPLUS

DOCUMENT NUMBER: 127:81793

TITLE: Preparation of hydroxyethylamine core structures as HIV and FIV protease inhibitors

INVENTOR(S): Wong, Chi-Huey; Snee, Deborah H.; Laslo, Karen

PATENT ASSIGNEE(S): Scripps Research Institute, USA; Wong, Chi-Huey; Snee, Deborah H.; Laslo, Karen

SOURCE: PCT Int. Appl., 202 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9721100	A1	19970612	WO 1996-US19571	19961209
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
CA 2238337	A1	19970612	CA 1996-2238337	19961209
AU 9712844	A	19970627	AU 1997-12844	19961209

10/580,011

AU 728373 B2 20010111
 EP 873519 A1 19981028 EP 1996-943657 19961209
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,
 IE, SI, LT, LV, FI, RO
 JP 2000502332 T 20000229 JP 1997-521485 19961209
 US 6900238 B1 20050531 US 1998-77712 19961209
 PRIORITY APPLN. INFO.: US 1995-568532 A2 19951207
 WO 1996-US19571 W 19961209
 OTHER SOURCE(S): MARPAT 127:81793
 GI



AB Combinatorial libraries of HIV and FIV protease inhibitors are characterized by α -keto amide or hydroxyethylamine core structures I and II [$n = 1, 2$; $R =$ one or more groups CONHMe_3 , CH_2OH , CH_2OMe , $\text{CH}_2\text{OCH}_2\text{Ph}$, OH , OCH_2Ph , C1-4 alkoxy, optionally nitro-substituted 2-, 3-, or 4-MeOC₆H₄CH₂O, 2,3- or 3,4-methylenedioxyphenylmethoxy, etc.; $R_1 =$ PhCH₂O₂C (Cbz), Me₃CO₂C (Boc), acyl; $R_2 =$ H, HO, PhCH₂O, C1-4 alkoxy, optionally nitro-substituted 2-, 3-, or 4-MeOC₆H₄CH₂O, 2,3- or 3,4-methylenedioxyphenylmethoxy] flanked by on one side by substituted pyrrolidines, piperidines, or azasugars and on the other side by Phe, Tyr, or substituted tyrosines. The libraries are synthesized via coupling of the nitrogen heterocycles with hydroxy acids, e.g. III, followed by oxidation to the keto amide, or a one-step coupling with epoxides, e.g. IV. Highly efficacious drug candidates are identified by screening the libraries for binding and inhibitory activity against both HIV and FIV protease. Drug candidates displaying clin. useful activity against both HIV and FIV protease are identified as being potentially resistive against a loss of inhibitory activity due to development of resistant strains of HIV.

IT 191850-51-8P 191850-64-3P 191850-67-6P
 191850-75-6P 191850-79-0P 191850-82-5P
 191850-85-8P 191850-88-1P

RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(preparation of hydroxyethylamine core structures as HIV and FIV protease inhibitors)

RN 191850-51-8 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3R,4R,5S)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

10/580,011


Absolute stereochemistry.



RN 191850-64-3 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3S,4R,5S)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-67-6 CAPLUS

CN Carbamic acid, [(1S)-3-[(2S,3R,4R,5R)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 191850-75-6 CAPLUS


CN Carbamic acid, [(1S)-3-[(2S,3S,4R,5R)-2-[[[(1,1-dimethylethyl)amino]carbonyl]-3,4,5-trihydroxy-1-piperidinyl]-2,3-dioxo-1-(phenylmethyl)propyl]-, phenylmethyl ester (9CI) (CA INDEX NAME)

10/580,011

RN 191850-42-7 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3S,4R,5S) - (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-45-0 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3R,4R,5R) - (CA INDEX NAME)


Absolute stereochemistry.



RN 191850-48-3 CAPLUS

CN 2-Piperidinecarboxamide, N-(1,1-dimethylethyl)-3,4,5-trihydroxy-,
(2S,3S,4R,5R) - (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 44 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1997:16443 CAPLUS

DOCUMENT NUMBER: 126:144017

TITLE: UK-2A, B, C and D, novel antifungal antibiotics from
Streptomyces sp. 517-02. II. Structural elucidation

AUTHOR(S): Hanafi, Muhammad; Shibata, Kozo; Ueki, Masashi;
Taniguchi, Makoto

CORPORATE SOURCE: Fac. Sci., Osaka City Univ., Osaka, 558, Japan

SOURCE: Journal of Antibiotics (1996), 49(12), 1226-1231

CODEN: JANTAJ; ISSN: 0021-8820

PUBLISHER: Japan Antibiotics Research Association

DOCUMENT TYPE: Journal

LANGUAGE: English

AB UK-2A, UK-2B, UK-2C and UK-2D, novel antibiotics produced by Streptomyces
sp. 517-02, exhibit strong antifungal activity. The structures were
elucidated based on spectral and chemical evidence that these compds. are the
derivs. of the nine-membered dilactone formed from serine and

10/580,011

4-hydroxypentanoic acid moiety.


IT 166820-04-8P 166820-06-0P

RL: SPN (Synthetic preparation); PREP (Preparation)
(structural elucidation of UK-2A, UK-2B, UK-2C and UK-2D, novel
antifungal antibiotics from Streptomyces sp. 517-02)

RN 166820-04-8 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, methyl ester
(CA INDEX NAME)


Absolute stereochemistry.



RN 166820-06-0 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]- (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L4 ANSWER 45 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1995:938109 CAPLUS

DOCUMENT NUMBER: 123:340945

TITLE: Preparation of pyridylcarbonylglycines and related
compounds as prolyl-4-hydroxylase inhibitors.

INVENTOR(S): Weidmann, klaus; Baringhaus, karl-Heinz; Tschank,
Georg; Bickel, Martin

PATENT ASSIGNEE(S): Hoechst A.-G., Germany

SOURCE: Eur. Pat. Appl., 35 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 661269	A1	19950705	EP 1994-120766	19941227
EP 661269	B1	19970326		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, NL, PT, SE				
DE 4344958	A1	19950706	DE 1993-4344958	19931230
DE 4439935	A1	19960515	DE 1994-4439935	19941109
CA 2138929	A1	19950701	CA 1994-2138929	19941222
TW 406076	B	20000921	TW 1994-83112023	19941222
AT 150749	T	19970415	AT 1994-120766	19941227
ES 2102132	T3	19970716	ES 1994-120766	19941227

10/580,011


ZA 9410340	A	19950831	ZA 1994-10340	19941228
JP 07242635	A	19950919	JP 1994-326903	19941228
US 5620995	A	19970415	US 1994-365411	19941228
NO 9405084	A	19950703	NO 1994-5084	19941229
AU 9481790	A	19950706	AU 1994-81790	19941229
AU 697015	B2	19980924		
CN 1126203	A	19960710	CN 1994-113548	19941230
PRIORITY APPLN. INFO.:			DE 1993-4344958	A 19931230
			DE 1994-4439935	A 19941109
OTHER SOURCE(S):		CASREACT 123:340945; MARPAT 123:340945		
GI				



AB Title compds. [I; X, Q = O, S; Y = N, CR3; m = 0,1; A = (substituted) alkylene; B = CO2H, NHSO2CF3, tetrazolyl, imidazolyl, 3-hydroxyisoxazolyl, etc.; R1-R3 = H, OH, halo, cyano, CF3, NO2, CO2H, alkyl, cycloalkyl, cycloalkoxy, aryl, aralkynyl, alkynylcarbonyl, alkylcarbonyloxy, carbamoyl, alkynyloxyalkyl, alkenyloxy, alkoxyalkoxy, alkynyl, retinyloxycarbonyl, alkenyloxycarbonyloxy, etc.; R1R2 or R2R3 = (CH2)_o in which 1-2 CH2 groups of the saturated or C:C unsatd. chain may be replaced by O, S, SO, SO2, imino, etc.; o = 3-5; R4 = H], were prepared for treatment of fibrotic disease (no data). Thus, 3-benzyloxypyridine-2-carboxylic acid (preparation given) in THF was treated sequentially with Et3N, pivaloyl chloride, and glycine Me ester hydrochloride at 0-20° to give 3-benzyloxypyridine-2-carboxylic acid (glycylmethyl ester)amide. This was hydrogenolyzed followed by saponification to give 3-hydroxypyridine-2-carboxylic acid glycylamide.

IT 170689-48-2P
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (preparation of pyridylcarbonylglycines and related compds. as prolyl-4-hydroxylase inhibitors)

RN 170689-48-2 CAPLUS
 CN 2-Pyridinecarboxamide, N-(2-amino-2-oxoethyl)-3-hydroxy-4-methoxy- (CA INDEX NAME)




IT 170689-59-5P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (preparation of pyridylcarbonylglycines and related compds. as prolyl-4-hydroxylase inhibitors)

10/580,011

RN 170689-59-5 CAPLUS

CN Glycine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, ethyl ester (CA INDEX NAME)



L4 ANSWER 46 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1995:671786 CAPLUS

DOCUMENT NUMBER: 123:164736

TITLE: The structures of UK-1 and UK-2, novel antibiotics from *Streptomyces* sp. 517-02

AUTHOR(S): Hanafi, O Muhammad; Kozo, Shibata; Masaru, Kashiwada; Masashi, Ueki; Makoto, Taniguchi

CORPORATE SOURCE: Faculty Science, Osaka City University, Japan

SOURCE: Tennen Yuki Kagobutsu Toronkai Koen Yoshishu (1994), 36th, 728-35
CODEN: TYKYDS

PUBLISHER: Nippon Kagakkai

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB The mycelial cake was extracted with acetone, and filtered. The filtrate was concentrated to give aqueous solution, which was extracted with chloroform.

Organic layer was concentrated to yield an oily material, followed by purification on silica gel column

chromatog. to give crude UK-1 and UK-2. Finally, the recrystn. of each fractions from MeOH, afforded UK-1 and UK-2. UK-1 (I), a novel metabolite, demonstrated potent cytotoxic activity against B16, Hela and P388 cells, and UK-2, novel complex of antibiotics, exhibited strong antifungal activity. Methylation of UK-1 by CH₃I and anhydrous K₂CO₃ in dry acetone gave monomethyl ether derivative, Me-UK-1. Alkaline hydrolysis of UK-1 afforded carboxylic acid derivative, DeMe-UK-1. Partial structures, A, B, and C were elucidated by COSY, and COLOC expts. Based on these results, the structure of UK-1 was deduced to be a novel benzoxazole dimer derivative UK-2, novel metabolite containing complex of antibiotics with strong antifungal activity, was purified by reverse phase HPLC, to give UK-2A, B, C and D. From NMR and mass spectral data, the structures of UK-2A, B, C and D were established as isobutyrate, tiglate, isovalerate, and 2-methylbutyrate of nine membered dilactone skeleton, resp. Based on the result of synthesis of hydrolysis products, the absolute configuration of UK-2 was identified.

IT 166820-04-8 166820-06-0

RL: MSC (Miscellaneous)


(structures of UK-1 and UK-2, novel antibiotics from *Streptomyces* sp. 517-02)

RN 166820-04-8 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl]-, methyl ester (CA INDEX NAME)

Absolute stereochemistry.


10/580,011



RN 166820-06-0 CAPLUS

CN L-Serine, N-[(3-hydroxy-4-methoxy-2-pyridinyl)carbonyl] - (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 47 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1993:6795 CAPLUS

DOCUMENT NUMBER: 118:6795

TITLE: Preparation of aztreonam hydrazides as antibiotics

INVENTOR(S): Treuner, Uwe D.

PATENT ASSIGNEE(S): E. R. Squibb and Sons, Inc., USA

SOURCE: U.S., 29 pp.

CODEN: USXXAM




DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5112968	A	19920512	US 1989-386070	19890728
PRIORITY APPLN. INFO.:			US 1989-386070	19890728
OTHER SOURCE(S):	MARPAT	118:6795		
GI				




AB Title compds. [I; G = CR₂R₃CONR_NRCOR₆; R = H, Me; R₁ = (substituted) Ph, -heterocyclyl; R₂, R₃ = H, alkyl; R₂R₃ = atoms to complete a carbocyclic ring; R₄, R₅ = H, (cyclo)alkyl, alkenyl, Ph, heterocyclyl, etc.; R₆ = pyridonyl group Q; Y₁ = CO₂H, CONH₂, OH, alkoxy, CHO, halomethyl, etc.; 1 of Y₂, Y₃ = OH and the other = H] were prepared as antibiotics (no data). Thus, maltol was converted in 11 steps to QCONHNH₂.CF₃CO₂H (Y₁ = CH₂OH, Y₂ = OH, Y₃ = H) which was condensed with aztreonam to give title compound II.

IT 144399-80-4P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and reaction of, in preparation of antibiotics)

RN 144399-80-4 CAPLUS

CN Carbamic acid, [(1,4-dihydro-3-hydroxy-4-oxo-2-pyridinyl)carbonyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



L4 ANSWER 48 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1990:526581 CAPLUS

DOCUMENT NUMBER: 113:126581

TITLE: Use of 2-hydroxymethyl-3,4,5-trihydroxypiperidines as antiviral agents

INVENTOR(S): Boeshagen, Horst; Junge, Bodo; Kinast, Guenther; Schueller, Matthias; Stoltefuss, Juergen; Paessens, Arnold

PATENT ASSIGNEE(S): Bayer A.-G., Fed. Rep. Ger.

SOURCE: Eur. Pat. Appl., 18 pp.

CODEN: EPXXDW

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 315017	A2	19890510	EP 1988-117701	19881025
R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, NL, SE				
DE 3737523	A1	19890518	DE 1987-3737523	19871105
US 5051407	A	19910924	US 1988-259932	19881019
JP 01151593	A	19890614	JP 1988-273377	19881031
PRIORITY APPLN. INFO.:			DE 1987-3737523	A 19871105


OTHER SOURCE(S): MARPAT 113:126581

GI For diagram(s), see printed CA Issue.


AB Title compds. [I; R₁ = H and R₃ = (substituted) aliphatic, cycloaliph., or aromatic residue which may contain hetero atoms; or R₁ = (substituted) aliphatic, cycloaliph., aromatic, or heterocyclic residue and R₃ = H, OH, OR₁, SH, SR₁, (substituted) amino or aminomethyl, CO₂H, etc.; or R₁ = (substituted) phenoxyalkyl, phenylthioalkyl, etc. and R₃ = H], especially 1-deoxynojirimycin derivs., are useful as medical virucides, especially against retroviruses. Thus, N-ethyl-1-deoxynojirimycin at 10 µg/mL showed a 50% inhibitory effect on the cytopathic activity of visna virus on sheep fibroblasts.

10/580,011

IT 81117-54-6 81117-55-7
RL: BIOL (Biological study)
(as medical virucides)
RN 81117-54-6 CAPLUS
CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-N-(phenylmethyl)- (CA INDEX NAME)




RN 81117-55-7 CAPLUS
CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-1-methyl-N-(phenylmethyl)- (CA INDEX NAME)



L4 ANSWER 49 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 1990:406035 CAPLUS
DOCUMENT NUMBER: 113:6035
TITLE: Ammoniomethylcephemcarboxylates as antibacterial agents and their preparation
PATENT ASSIGNEE(S): Fujisawa Pharmaceutical Co., Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 16 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 02015090	A	19900118	JP 1989-133559	19890526
PRIORITY APPLN. INFO.:			GB 1988-13945	A 19880613
OTHER SOURCE(S):	MARPAT	113:6035		


GI



- AB The title compds. I [R1 = (protected) amino; R2 = organic group; R3,R4 = alkyl; R5,R6 = (protected) hydroxy; A = alkylene; X = NH, O; Q, Z = N, CH] and their pharmaceutically acceptable salts were prepared Reaction of 7β-[2-(5-amino-1,2,4-thiadiazol-3-yl)-2-(1-carboxy-1-methylethoxyimino)acetamido]-3-chloromethyl-3-cephem-4-carboxylic acid (syn isomer) CF₃CO₂H salt with N,N-dimethyl-2-(3,4-diacetoxybenzoyloxy)ethylamine, followed by deprotection, gave 7β-[2-(5-amino-1,2,4-thiadiazol-3-yl)-2-(1-carboxy-1-methylethoxyimino)acetamido]-3-[N,N-dimethyl-N-[2-(3,4-dihydroxybenzoyloxy)ethyl]ammoniomethyl]-3-cephem-4-carboxylate (syn isomer) (II). II in vitro exhibited a MIC of ≤0.025 μg against *Pseudomonas aeruginosa* 26.
- IT 127450-16-2P
RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and reaction of, in preparation of antibacterial agent)
- RN 127450-16-2 CAPLUS
- CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-3-methanaminium, 2-carboxy-N-[2-[[[(3,4-dihydroxy-2-pyridinyl)carbonyl]amino]ethyl]-7-[[[2-(formylamino)-4-thiazolyl](methoxyimino)acetyl]amino]-N,N-dimethyl-8-oxo-, chloride, hydrochloride, [6R-[6α,7β(Z)]]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

PAGE 1-A

● Cl⁻


PAGE 2-A

● x HCl

10/580,011

TITLE: Preparation of sugar lactams as antiinflammatories and pharmaceutical compositions containing them
INVENTOR(S): Tsuruoka, Takashi; Yuda, Yasukatsu; Nakabayashi, Akira; Katano, Kiyoaki; Sezaki, Masaji; Kondo, Shinichi
PATENT ASSIGNEE(S): Meiji Seika Kaisha, Ltd., Japan
SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.
CODEN: JKXXAF
DOCUMENT TYPE: Patent
LANGUAGE: Japanese.
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 63216867	A	19880909	JP 1987-50100	19870306
JP 06076379	B	19940928		
PRIORITY APPLN. INFO.:			JP 1987-50100	19870306
OTHER SOURCE(S):	MARPAT	111:78547		
GI				




AB The title compds. [I; Z = CH₂O-W, CO₂Y₁, CONHY₂; R = H, acyl, Bz; W = (substituted) phenylsulfonyl, aralkylsulfonyl, heterocyclylsulfonyl, (substituted) alkanoyl, (substituted) benzoyl, heterocyclylcarbonyl, CHR₁R₂; R₁, R₂ = H, alkyl, (substituted) Ph, (substituted) naphthyl, heterocyclyl; Y₁ = alkyl, aralkyl; Y₂ = alkyl, (substituted) Ph, aralkyl, heterocyclyl], useful as antiinflammatories, are prepared
D-Gluco-8-lactam was reacted with Ph₂CHCOCl in pyridine to give I (R = H, Z = CH₂O₂CCHPh₂) (II), which in the carrageenin test showed 59.2% inhibition of inflammation, vs. 38.2% for aspirin. A tablet containing II 50, lactose 280, potato starch 80, polyvinylpyrrolidone 11, and Mg stearate 5 mg was formulated.

IT 121715-76-2P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of, as antiinflammatory)

RN 121715-76-2 CAPLUS

CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-oxo-N-(phenylmethyl)-, [2S-(2 α ,3 β ,4 α ,5 β)]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 51 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1982:523505 CAPLUS

DOCUMENT NUMBER: 97:123505

ORIGINAL REFERENCE NO.: 97:20437a,20440a

TITLE: Structure of rubradirin

AUTHOR(S): Hoeksema, H.; Mizzak, S. A.; Baczynskyj, L.;
Pschigoda, L.

CORPORATE SOURCE: Res. Lab., Upjohn Co., Kalamazoo, MI, 49001, USA

SOURCE: Journal of the American Chemical Society (1982),
104(19), 5173-81

CODEN: JACSAT; ISSN: 0002-7863

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The antibiotic rubradirin from *Streptomyces achromogenes* consists of a central moiety, 3,4-dihydroxydipicolinic acid, of which the 2-carboxyl group is esterified by a large ansamycin-like moiety while the 6-carboxyl forms an amide with 3-amino-4-hydroxy-7-methoxycoumarin, a compound of the type found in the novobiocins. Position 4 is glycosylated with a nitro sugar, rubranitrose, which is epimeric with evernitrose, found in 3rd class of antibiotics, the everninomycins. Rubradirins B and C are members of the rubradirin complex which lack rubranitrose and also have slight modifications elsewhere.

IT 69282-24-2P 69282-25-3P 71502-31-3P

71502-32-4P 71502-33-5P 82537-38-0P


82537-39-1P

RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)

RN 69282-24-2 CAPLUS

CN 2,6-Pyridinedicarboxamide, N2-ethyl-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro-β-L-xylo-hexopyranosyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 69282-25-3 CAPLUS

CN 2,6-Pyridinedicarboxamide, 3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-

10/580,011


nitro- β -L-xyllo-hexopyranosyl)oxy] - (9CI) (CA INDEX NAME)

Absolute stereochemistry.




RN 71502-31-3 CAPLUS

CN 2-Pyridinecarboxylic acid, 1,4-dihydro-5-hydroxy-6-[(methylamino)carbonyl]-4-oxo- (CA INDEX NAME)




RN 71502-32-4 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-methyl-4-oxo- (CA INDEX NAME)



RN 71502-33-5 CAPLUS

CN 2,6-Pyridinedicarboxamide, 1,4-dihydro-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-oxo- (CA INDEX NAME)




RN 82537-38-0 CAPLUS

10/580,011


CN 2-Pyridinecarboxylic acid, 5-hydroxy-6-[(methyamino)carbonyl]-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro-β-L-arabino-hexopyranosyl)oxy]-, hydrazide (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 82537-39-1 CAPLUS

CN 2-Pyridinecarboxylic acid, 1,4-dihydro-5-hydroxy-6-[(methyamino)carbonyl]-4-oxo-, hydrazide (CA INDEX NAME)



L4 ANSWER 52 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1982:117597 CAPLUS

DOCUMENT NUMBER: 96:117597

ORIGINAL REFERENCE NO.: 96:19243a,19246a

TITLE: Herbicidal composition containing piperidine derivatives

INVENTOR(S): Berg, Dieter; Junge, Bodo; Stoltefuss, Juergen; Schmidt, Robert Rudolf

PATENT ASSIGNEE(S): Bayer A.-G. , Fed. Rep. Ger.

SOURCE: Ger. Offen., 30 pp.

CODEN: GWXXBX


DOCUMENT TYPE: Patent

LANGUAGE: German


FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 3024901	A1	19820128	DE 1980-3024901	19800701
PRIORITY APPLN. INFO.:			DE 1980-3024901	A 19800701
OTHER SOURCE(S):		CASREACT 96:117597		
GI				



- AB The 2-hydroxymethyl-3,4,5-trihydroxypiperidine derivs. I (R1 = H, alkyl, XR3 alkenyl, etc.; R2 = H, CN, OH, CH2OH, NHMe, etc.; R3 = aryl, aryloxy, arylmercapto, pyridyl, etc.; X = alkylene or alkenylene) are herbicides. Thus, in pot expts., N-(β -hydroxyethyl)-1-deoxynojirimycin [72432-03-2] (40 kg/ha) totally controlled *Lepidium*, *Portulaca*, and *Poa*. The synthesis of I is given.
- IT 81117-54-6P 81117-55-7P
 RL: AGR (Agricultural use); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses) (preparation of, as herbicide)
- RN 81117-54-6 CAPLUS
- CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-N-(phenylmethyl)- (CA INDEX NAME)



- RN 81117-55-7 CAPLUS
- CN 2-Piperidinecarboxamide, 3,4,5-trihydroxy-6-(hydroxymethyl)-1-methyl-N-(phenylmethyl)- (CA INDEX NAME)



L4 ANSWER 53 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1981:30737 CAPLUS

DOCUMENT NUMBER: 94:30737

ORIGINAL REFERENCE NO.: 94:5075a,5078a

TITLE: Oxazolo[4,5-c]coumarin derivative

INVENTOR(S): Hoeksema, Herman

PATENT ASSIGNEE(S): Upjohn Co., USA

SOURCE: U.S., 7 pp. Division of U.S. No. 4,137,410.
 CODEN: USXXAM

DOCUMENT TYPE: Patent

10/580,011


LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 3
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 4199507	A	19800422	US 1978-938688	19780831
US 4137410	A	19790130	US 1978-874767	19780206
JP 53137998	A	19781201	JP 1978-50259	19780428
GB 1576082	A	19801001	GB 1978-17205	19780502
GB 1576084	A	19801001	GB 1978-32822	19780502
GB 1576083	A	19801001	GB 1978-32823	19780502
GB 1576085	A	19801001	GB 1978-32824	19780502
GB 1576086	A	19801001	GB 1978-32825	19780502
GB 1576087	A	19801001	GB 1978-32826	19780502
US 4154939	A	19790515	US 1978-938602	19780831
US 4154940	A	19790515	US 1978-938687	19780831
US 4154938	A	19790515	US 1978-938689	19780831
US 4171437	A	19791016	US 1978-938603	19780831
US 4171436	A	19791016	US 1978-938686	19780831
US 4182855	A	19800108	US 1978-938606	19780831
US 4220786	A	19800902	US 1978-938685	19780831
FR 2411195	A1	19790706	FR 1979-4677	19790223
FR 2411195	B1	19810828		
FR 2411206	A1	19790706	FR 1979-4678	19790223
FR 2411206	B1	19810828		
FR 2411203	A1	19790706	FR 1979-4679	19790223
FR 2411203	B1	19830121		
FR 2411189	A1	19790706	FR 1979-4680	19790223
FR 2411189	B1	19810828		
FR 2411190	A1	19790706	FR 1979-4681	19790223
FR 2411190	B1	19821126		

PRIORITY APPLN. INFO.:

US 1977-793785 A2 19770505
 US 1978-874767 A3 19780206

GI



AB The fused compound I was obtained from rubradirin. A mixture of rubradirin, Ac₂O, and pyridine was refluxed 4 h to give I. The basic degradation of rubradirin gave rubransarol A, which showed bactericidal activity.

IT 69282-25-3P 75945-33-4P


RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

RN 69282-25-3 CAPLUS

CN 2,6-Pyridinedicarboxamide, 3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro-β-L-xylo-hexopyranosyl)oxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

10/580,011



RN 75945-33-4 CAPLUS

CN 2,6-Pyridinedicarboxamide, N2-ethyl-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-4-[(tetrahydro-5-methoxy-4,6-dimethyl-4-nitro-2H-pyran-2-yl)oxy]- (9CI) (CA INDEX NAME)



L4 ANSWER 54 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1980:76522 CAPLUS

DOCUMENT NUMBER: 92:76522

ORIGINAL REFERENCE NO.: 92:12611a,12614a

TITLE: Degradation of rubradirin and its B form

INVENTOR(S): Hoeksema, Herman

PATENT ASSIGNEE(S): Upjohn Co., USA

SOURCE: Fr. Demande, 22 pp.

CODEN: FRXXBL

DOCUMENT TYPE: Patent

LANGUAGE: French

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
FR 2409999	A1	19790622	FR 1978-13250	19780503
FR 2409999	B1	19830114		
US 4137410	A	19790130	US 1978-874767	19780206
JP 53137998	A	19781201	JP 1978-50259	19780428
GB 1576082	A	19801001	GB 1978-17205	19780502
GB 1576084	A	19801001	GB 1978-32822	19780502
GB 1576083	A	19801001	GB 1978-32823	19780502
GB 1576085	A	19801001	GB 1978-32824	19780502
GB 1576086	A	19801001	GB 1978-32825	19780502

10/580,011


GB 1576087	A	19801001	GB 1978-32826	19780502
US 4154939	A	19790515	US 1978-938602	19780831
US 4154940	A	19790515	US 1978-938687	19780831
US 4154938	A	19790515	US 1978-938689	19780831
US 4171437	A	19791016	US 1978-938603	19780831
US 4171436	A	19791016	US 1978-938686	19780831
US 4182855	A	19800108	US 1978-938606	19780831
US 4220786	A	19800902	US 1978-938685	19780831
FR 2411195	A1	19790706	FR 1979-4677	19790223
FR 2411195	B1	19810828		
FR 2411206	A1	19790706	FR 1979-4678	19790223
FR 2411206	B1	19810828		
FR 2411203	A1	19790706	FR 1979-4679	19790223
FR 2411203	B1	19830121		
FR 2411189	A1	19790706	FR 1979-4680	19790223
FR 2411189	B1	19810828		
FR 2411190	A1	19790706	FR 1979-4681	19790223
FR 2411190	B1	19821126		

PRIORITY APPLN. INFO.:

US 1977-793785	A	19770505
US 1978-874767	A	19780206

OTHER SOURCE(S): MARPAT 92:76522

GI.



I

AB The rubradirin aglycone I was obtained by the acidic degradation of rubradirin; I showed bactericidal activity. A mixture of rubradirin, HOAc, and water was stirred 6 days at room temperature to give I and L- and D-rubranitrose. The basic degradation of rubradirin gave rubransarol A and a rubradiric acid, and the former exhibited bactericidal activity. An isomer of rubransarol A and a rubradiric acid aglycone were obtained from rubradirin B.


IT 69282-24-2P 69282-25-3P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of)

RN 69282-24-2 CAPLUS
 CN 2,6-Pyridinedicarboxamide, N2-ethyl-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-

10/580,011

oxo-2H-1-benzopyran-3-yl)-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro- β -L-xylo-hexopyranosyl)oxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN. 69282-25-3 CAPLUS

CN 2,6-Pyridinedicarboxamide, 3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro- β -L-xylo-hexopyranosyl)oxy]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 55 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1979:575314 CAPLUS

DOCUMENT NUMBER: 91:175314

ORIGINAL REFERENCE NO.: 91:28283a,28286a

TITLE: The chemistry of rubradirin. III. The rubradiric acids and the structure of rubradirin

AUTHOR(S): Hoeksema, Herman; Mizzak, Stephen A.; Baczynskyj, Lubomir

CORPORATE SOURCE: Pharm. Res. Dev., Upjohn Co., Kalamazoo, MI, 49001, USA

SOURCE: Journal of Antibiotics (1979), 32(7), 773-6
CODEN: JANTAJ; ISSN: 0021-8820

DOCUMENT TYPE: Journal

LANGUAGE: English

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB Basic hydrolysis of rubradirin (I) and rubradirin B (II) gave in addition to rubransarols A and B, rubradiric acid A (III) and rubraduric acid B.

IT 71502-31-3P 71502-32-4P 71502-33-5P


10/580,011

RL: PREP (Preparation)

(isolation of, in structure proof of rubradiric acids and rubradirin)


RN 71502-31-3 CAPLUS

CN 2-Pyridinecarboxylic acid, 1,4-dihydro-5-hydroxy-6-[(methylamino)carbonyl]-4-oxo- (CA INDEX NAME)




RN 71502-32-4 CAPLUS

CN 2-Pyridinecarboxamide, 1,4-dihydro-3-hydroxy-N-methyl-4-oxo- (CA INDEX NAME)



RN 71502-33-5 CAPLUS

CN 2,6-Pyridinedicarboxamide, 1,4-dihydro-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-oxo- (CA INDEX NAME)



L4 ANSWER 56 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1979:103973 CAPLUS

DOCUMENT NUMBER: 90:103973

ORIGINAL REFERENCE NO.: 90:16427a,16430a

TITLE: Decomposition products of antibiotics rubradirin and rubradirin B

INVENTOR(S): Hoeksema, Herman

PATENT ASSIGNEE(S): Upjohn Co., USA

SOURCE: Ger. Offen., 35 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent


LANGUAGE: German

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 2816052	A1	19781116	DE 1978-2816052	19780413
US 4137410	A	19790130	US 1978-874767	19780206
JP 53137998	A	19781201	JP 1978-50259	19780428
GB 1576082	A	19801001	GB 1978-17205	19780502
GB 1576084	A	19801001	GB 1978-32822	19780502
GB 1576083	A	19801001	GB 1978-32823	19780502
GB 1576085	A	19801001	GB 1978-32824	19780502
GB 1576086	A	19801001	GB 1978-32825	19780502
GB 1576087	A	19801001	GB 1978-32826	19780502
US 4154939	A	19790515	US 1978-938602	19780831
US 4154940	A	19790515	US 1978-938687	19780831
US 4154938	A	19790515	US 1978-938689	19780831
US 4171437	A	19791016	US 1978-938603	19780831
US 4171436	A	19791016	US 1978-938686	19780831
US 4182855	A	19800108	US 1978-938606	19780831
US 4220786	A	19800902	US 1978-938685	19780831
FR 2411195	A1	19790706	FR 1979-4677	19790223
FR 2411195	B1	19810828		
FR 2411206	A1	19790706	FR 1979-4678	19790223
FR 2411206	B1	19810828		
FR 2411203	A1	19790706	FR 1979-4679	19790223
FR 2411203	B1	19830121		
FR 2411189	A1	19790706	FR 1979-4680	19790223
FR 2411189	B1	19810828		
FR 2411190	A1	19790706	FR 1979-4681	19790223
FR 2411190	B1	19821126		
PRIORITY APPLN. INFO.:			US 1977-793785	A 19770505
			US 1978-874767	A 19780206

GI



AB The acidic and alkaline decomposition of antibiotics rubradirin and rubradirin
 B gave 8 identified compds. including I and a configurational isomer of I.
 IT 69282-24-2P 69282-25-3P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation of, from rubradirin)
 RN 69282-24-2 CAPLUS
 CN 2,6-Pyridinedicarboxamide, N2-ethyl-3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro-β-L-xylo-hexopyranosyl)oxy]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



RN 69282-25-3 CAPLUS

CN 2,6-Pyridinedicarboxamide, 3-hydroxy-N6-(4-hydroxy-7-methoxy-2-oxo-2H-1-benzopyran-3-yl)-N2-methyl-4-[(2,3,6-trideoxy-3-C-methyl-4-O-methyl-3-nitro- β -L-xylo-hexopyranosyl)oxy]-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 57 OF 57 CAPLUS COPYRIGHT 2008 ACS on STN

ACCESSION NUMBER: 1952:67070 CAPLUS

DOCUMENT NUMBER: 46:67070

ORIGINAL REFERENCE NO.: 46:11209f-g

TITLE: 4 - Pyronecarboxylic acids and their transformations

AUTHOR(S): Belonosov, I. S.

CORPORATE SOURCE: Khabarovsk Med. Inst.

SOURCE: Zhurnal Prikladnoi Khimii (Sankt-Peterburg, Russian

Federation) (1951), 24, 113-16

CODEN: ZPKHAB; ISSN: 0044-4618

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB Meconic acid (50 g.) in 150 ml. concentrated NH_4OH , evaporated, and then boiled until CO_2 evolution ceased, gave the NH_4 salt of comenamic acid (3,4-dihydroxypicolinic acid); treatment with 10% HCl yielded 34.6% free acid, m. $260-2^\circ$, which with EtOH and dry HCl with cooling gave 40% Et ester, m. $204-5^\circ$. This (25 g.) let stand overnight with 125 ml. Et_2NH , filtered, and the solid dried, taken up in a little absolute EtOH , saturated with HCl , and diluted with Et_2O , gave a precipitate of crude N,N-diethyl-3,4-dihydroxypicolinamide- HCl ; purified by repeated treatment with $\text{EtOH-Et}_2\text{O}$, it m. 159° (13.3% yield). Comenic acid with dry HCl in EtOH gave 40.6% Et ester, m. 127° , which with Et_2NH as above gave 49.5% diethylamine- HCl , m. 168° , decompose slowly on standing in the open air; it is toxic to the isolated frog heart.


IT 856834-25-8P, Picolinamide, N,N-diethyl-3,4-dihydroxy-, hydrochloride

RL: PREP (Preparation)
(preparation of)

10/580,011

RN 856834-25-8 CAPLUS

CN Picolinamide, N,N-diethyl-3,4-dihydroxy-, hydrochloride (5CI) (CA INDEX NAME)



● HCl

=>